

Baseline Assessments and Analysis of Fish, Macroinvertebrates and Herpetofauna in the Otter Creek Coal Tracts Area of Powder River County

Prepared for:

Pat Farmer, Project Manager/Director
Westech Environmental Services
Helena Field Office



Prepared by:

David Stagliano, Aquatic Ecologist
Montana Natural Heritage Program
a cooperative program of the
Montana State Library and the University of Montana

January 2012



Executive Summary

We summarize the first year of baseline surveys for the Aquatic Assessment of Fish, Macroinvertebrates, and Herpetofauna in the Otter Creek coal tracts area. Project goals were: 1) to **conduct standardized surveys** and collect baseline information on the aquatic and riparian communities occurring in Otter Creek and three tributaries (seasonally and spatially oriented) prior to coal development, 2) to **assess aquatic community integrity** and condition by interpreting key indicators recorded at sites using standardized protocols and biotic thresholds, and to compare these against reference condition standards. These data collected represent pre-development (i.e. pre-impact, BACI design) conditions at the local reach scale.

Habitat assessments, herpetofauna, macroinvertebrate and fish surveys were performed seasonally at three predetermined mainstem Otter Creek reaches (control, impact and downstream) and three tributaries coinciding with established surface water quality stations during 2011. Additionally, we added fish sampling visits to Otter Creek Impact #2 (upstream of the Threemile Creek confluence) because Threemile Creek remained “dry” during all visits. In total, we performed 15 surveys for fish during the visits: 11 at four mainstem Otter Creek reaches and four surveys at two tributary streams. Fifteen macroinvertebrate samples were collected during the visits; neither taxa survey was conducted at Threemile Creek in any season due to lack of surface water present. All six *a priori* stream reaches were visually surveyed for amphibians or reptiles during all visits. Biological community integrity was calculated for 15 fish surveys using Fish Integrated Biotic Indices (IBI's) and Observed/Expected Models (O/E), while the 15 macroinvertebrate samples were assessed with Montana DEQ's multi-metric indices (MT MMI).

Habitat Evaluations. Of the seven sampling reaches evaluated in the study area, we found three in Proper Functioning Condition (PFC) with a stable trend and four were Functional at Risk (FAR). Reasons that sites ranked FAR were likely due to anthropogenic habitat alteration by cattle (Home Creek {Otter_1A} and Threemile Creek {Otter_3m}) or stream manipulation (Otter Creek JT-Trussler and Otter Creek #16). Highest site integrity scores using both the BLM Habitat and PFC Assessment methods were recorded at the Otter Creek sites #23 (Tenmile Creek) and #22 (control-Denson reach). Sites with lower habitat scores were structurally degraded predominately by cattle use and had high associated Livestock Use Indices (LUI) (Home Creek, Threemile and Otter Creek #16-fall). Point conductivity measurements recorded at all Otter Creek mainstem sites across most seasons were above the threshold for impairment levels ($>500\mu\text{s}$, DEQ 2006), and Home Creek site 1A had visible signs of natural gas seepage from the sediments.

Macroinvertebrate Communities: Overall, 104 unique macroinvertebrate taxa were reported from the 15 macroinvertebrate assessment samples. One known Montana species of concern (SOC), the mayfly, *Caenis youngi* was collected in fair numbers at the control site, Otter Creek #22. Average macroinvertebrate taxa richness per site was 29.5 and the highest taxa richness was 41 taxa reported at the Otter Creek JT site. Using the Montana DEQ macroinvertebrate multimetric index (MMI), four of the five sites (12 of 15 samples) were ranked non-impaired (good to excellent biological integrity), while all three samples from Tenmile Creek were ranked marginally impaired. Stream sites that maintained flowing, connected water scored higher with the MMI than sites with

interrupted pool areas. Overall, mainstem sites evaluated in the Otter Creek study received significantly higher macroinvertebrate MMI scores than those in the tributaries (T-test, $p < 0.01$). MMI's did not significantly differ on Otter Creek mainstem Pre-Impact Control, Impact or Downstream Sites (T-test, $p > 0.05$), despite the fish communities reflecting a decrease in biotic integrity.

Fish Communities. Overall, nine fish species (five native/four introduced) were identified from 37,679 individuals collected from 15 site visits (Table 4). One potential species of concern (PSOC), the brassy minnow, was collected at five of six sites. Average fish species per Otter Creek mainstem site across all seasons was $6.5 (\pm 0.8 \text{ SE})$, while the tributary sites averaged 1.75 species (Table 4). All fish presence sites also reported at least one species of amphibian. Lake chubs had the highest site occupancy rate at 93% (14 of 15 visits) followed by fathead and brassy minnows at 80% (12 of 15 visits). Fish data collected in previous years from three sites within the study area showed similar biological integrity over time. The most diverse site in the study area was Otter Creek JT site with nine species, while the most intact sites were Otter Creek Site #22 at Denson's (four native) and the Tenmile Creek spring survey (two native spp.). Using Montana's Prairie Fish IBI, 9 of the 15 fish sites were ranked non-impaired (good biological integrity), two were slightly impaired (moderate integrity) and four were moderately impaired (poor biotic integrity). Fish IBI's decreased going downstream on Otter Creek, and the Pre-Impact Control Site scored significantly higher than Impact or Downstream sites (T-test, $p < 0.05$).

Amphibian and Reptile Incidentals. Eight herpetofauna species were observed, collected in dipnets/seines or incidentally recorded in conjunction with the fish, habitat and macroinvertebrate surveys. We reported four amphibian species, of which, Woodhouse's Toad (*Bufo woodhousii*) had the highest site occupancy, occurring at six of seven sites, followed by the Northern Leopard Frog (*Rana pipiens*) and Tiger Salamander (*Ambystoma tigrinum*) recorded at five and four sites, respectively. Boreal Chorus Frog (*Pseudacris maculata*) was detected at three sites during the spring visits only. We also recorded four reptile species (in order of site occurrence): Western Rattlesnake (*Crotalus viridis*), Painted Turtle (*Chrysemys picta*), Terrestrial Garter Snake, (*Thamnophis elegans*) and Snapping Turtle (*Chelydra serpentina*) (SOC).

Conclusions. The aquatic community sampling sites chosen for pre-impact baseline data were representative of the range of stream classes found in the Otter Coal Tracts project area: Ephemeral, Intermittent and Perennial Prairie Streams. Despite this having been an unusually high water year for the region, stream communities that we sampled encompassed the range of expected species to occur in these stream types. Biotic integrity of sites was initially higher in the upstream control reaches of Otter Creek. One extraordinary finding was the high density and large biomass of fish inhabiting the stream reach below Trusler's Ranch road crossing ($\sim 20,000$ fish in the 300 m reach fall survey, 15 fish/m^2); this was most likely an artifact of the impassible culvert located here and the fish "stacking up" downstream of this barrier. Density dependent fish anomalies (lesions and parasites, i.e., yellow grub and anchorworm) were prevalent at this site, indicating the fish were likely experiencing stress from overcrowding. Benthic macroinvertebrate densities at this site were low, but, more diverse than other sites, indicating that the fish may have been consuming large portions of the available insect production, having a top-down community effect.

TABLE OF CONTENTS

INTRODUCTION	6
METHODS	
<i>Study Area</i>	6
<i>Habitat Assessments</i>	7
<i>Macroinvertebrate Community Surveys</i>	9
<i>Fish and Amphibian Surveys Surveys</i>	10
RESULTS	
<i>Habitat Assessments</i>	11
<i>Macroinvertebrate Community Surveys</i>	12
<i>Fish Communities</i>	14
<i>Hepetofauna Communities</i>	18
CONCLUSIONS	19
SITE PHOTOS.....	21
LITERATURE CITED.....	25

APPENDIX A. Fish data and IBI metric calculations collected from Otter Creek Project Sites

APPENDIX B. Macroinvertebate taxa list, abundance and metrics for the 15 collection sites

List of Figures

Figure 1. USGS gage data for Otter Creek at Ashland in 2011	8
Figure 2. Macroinvertebrate D-net Sampling procedure.....	9
Figure 3. Brassy minnows and lake chubs in the seine at Otter 22	10
Figure 4. Photo of a structurally cattle-damaged, functional-at-risk (FAR) stream reach.....	11
Figure 5. DEQ MMI scores by site and season	13
Figure 6. Average macroinvertebrate MMI scores by site type.....	14
Figure 7. Fish IBI seasonality recorded during 2011 surveys	16
Figure 8. Comparison of the Fish IBI and O/E scores between sites and season	17
Figure 9. Average fish IBI scores by BACI design site type	17

List of Tables

Table 1. Otter Creek Study site GPS locations of reach	7
Table 2. Impairment determinations from the MMI and O/E (RIVPACS) models	9
Table 3. Fish metrics and classification of fish species captured in the study area.....	11
Table 4. Macroinvertebrate sample characteristics and metrics used for the DEQ MMI	13
Table 5. Fish abundance, IBI and O/E scores for the 12 surveys reporting fish.....	15
Table 6. Species of vertebrates collected during the aquatic surveys in 2011.....	18
Table 7. Otter Creek Temporal Habitat Scores, Fish IBIs and MMIs	19

Acknowledgements

We would like to thank Pat Farmer of Westech Environmental Services of Helena for funding this project under agreement # L08AC13222. Report review and editing was provided by Stephen Fernandes and Linda Vance. Field work coordination, landowner contacts and logistics were greatly expediated by Carl Lenz and Heidi Kaiser (Hydrometrics, Inc.). Fieldwork assistance provided by Stephen Fernandes (MTNHP) was invaluable.

All photos in the report were taken by MTNHP personnel, unless otherwise noted

Introduction

The Otter Creek basin in southeast Montana is currently undergoing exploration for a large, open-pit coal mine. Baseline data on the condition of the ecosystem prior to coal mine development (pre-impact) is essential to determine what effects the coal extraction might have on the fish and wildlife in, and downstream of, the effected extraction area (post-impact). The initial mining rights transfer determined that no Federally Listed species under the ESA would be affected in the coal tracts area (BLM EA 2002). But this EA did not take into account the presence of Montana species of concern (SOC) or other ecologically sensitive native species assemblages. Despite numerous projects undertaken to document and monitor biological communities in the Powder and Tongue River watersheds in the wake of CBNG development (Confluence Consulting Inc. 2003, 2004; Stagliano 2006; Davis et al. 2009; Maxell 2009; Petersen et al. 2009; Senecal 2009, Stagliano 2011), large gaps still exist in basic baseline surveys for riparian macroinvertebrate, fish, and herpetofauna in the Otter Creek basin. Fish communities have been documented to be seasonally variable in prairie streams (Bramblett and Fausch 1991), thus sampling across all seasons is a good strategy to document baseline community differences. Many of the remaining data gaps involve small prairie streams that constitute the Northwestern Great Plains Intermittent Prairie Stream ecological system (Stagliano 2005). This stream system may have downstream connectivity early in the season for potential fish spawning and nursery areas (Smith and Hubert 1989, Bramblett 2000) or no fish colonization at all, but by summer often becomes a string of isolated pools that are important breeding and rearing areas for amphibians (Stagliano 2011). Identifying baseline communities and conditions in streams of the coal tracts area (i.e., the presence of fish, macroinvertebrate, and herpetofauna) prior to coal development is essential to understanding and potentially mitigating impacts to habitats and species during and after coal extraction.

Methods

Study Sites

The aquatic community sampling sites chosen for pre-impact baseline data are representative of the range of streams found in the Otter Coal Tracts project area: Ephemeral, Intermittent and Perennial Prairie Streams. Three predetermined mainstem Otter Creek reaches (control, impact and downstream) and three tributaries coinciding with established surface water quality stations were visited seasonally (May, July, October) during 2011 (Table 1). Additionally, we added fish sampling visits to Otter Creek site #2 near the Threemile Creek confluence, since Threemile Creek remained “dry” during all visits (see Site Photos). Study site visits were timed with season and discharge (Figure 1); although we tried to coordinate field visits with baseflow water levels, 2011 was an


Table 1. Otter Creek Coal Study Site GPS Locations at the top (T) and bottom (B) of the assessment reach. Aquatic Ecological System (AES) code described in text followed Stagliano (2005).

Site Code	Site Name	Type	AES code	Latitude	Longitude	Elev (m)	Comment
Otter_23	Tenmile Creek (T)	Control	D005/E005	45.43409	-106.13403	961	Spring survey for fish above the road reach, dry during July, Oct. visits
Otter_23	Tenmile Creek (B)	Control	D005/E005	45.43465	-106.13253	958	
Otter_22	Otter Creek 22 (T)	Control	D005/C005	45.43035	-106.14428	951	Top of reach began ~60m below Tenmile Creek road, proceeded 300m downstream
Otter_22	Otter Creek 22 (B)	Control	D005/C005	45.43274	-106.14366	948	
Otter_16	Otter Creek 16 (T)	Impact	C005	45.48514	-106.16487	938	Top of reach began ~60m above the stream crossing and went 240m below
Otter_16	Otter Creek 16 (B)	Impact	C005	45.48365	-106.16725	937	
Otter_3m	Threemile Creek (T)	Impact	E005	45.51054	-106.16288	933	Spring survey contained standing water in cattle hoof prints, dry during July, Oct. visits
Otter_3m	Threemile Creek (B)	Impact	E005	45.50955	-106.16960	928	
Otter_2	Otter Creek 2 (T)	Impact	C005	45.50475	-106.17493	929	Site added for fish surveys during July visit (unseizable in Spring), downstream of road crossing
Otter_2	Otter Creek 2 (B)	Impact	C005	45.50561	-106.17561	928	
Otter_JT	Otter Creek JT (T)	Down	C005	45.55675	-106.21798	910	Top of reach began ~80m below the ranch road/impassable culvert, proceeded 300m down
Otter_JT	Otter Creek JT (B)	Down	C005	45.55782	-106.21770	909	
Otter_1A	Home Creek (B)	Down	D005/E005	45.54483	-106.18717	952	Bottom of reach began ~500m above the Otter Creek road crossing, proceeded 300m upstream
Otter_1A	Home Creek (T)	Down	D005/E005	45.54422	-106.18947	950	

extreme high water year (average yearly discharge 14.9 vs. 3.96 cfs in 2010). Therefore, we had to take advantage of sampling conditions as they presented themselves, sometimes not during the most ideal sampling periods (e.g., May visit @ 19 cfs between two high water pulses, Figure 1). Summer and fall visits were closer to base flow at 11 and 7 cfs, respectively. Habitat assessments, herpotofauna, macroinvertebrate and fish surveys were performed during the same site visit. In total, we surveyed 15 reaches for fish during the visits: four mainstem Otter Creek reaches (11 surveys) and two tributary streams (4 surveys). Fifteen macroinvertebrate samples were collected during the visits; neither survey was conducted at Threemile Creek in any season due to lack of surface water present. Spring and fall macroinvertebrate samples were collected outside of the MTDEQ recommend sampling time frame (June 1st-September 15th) (MTDEQ 2006), but this period

was largely derived for mountain streams. All six a priori stream reaches were visually surveyed for amphibians or reptiles during all visits.

Figure 1. Discharge reported at the USGS gage in Ashland, MT. Arrows indicate sampling dates.



Riparian Assessments

The assessment stream reach was divided into 10 equally spaced transects according to BLM and EMAP protocols (BLM 2008b; Lazorchak et al. 1998). The downstream transect was marked (GPS, flagging and photo point) as the bottom of the reach and all ecological assessment protocols started from this point and continued upstream for 300m (designated the assessment area or “AA”) to the top of the reach which was also marked. Parameters recorded at each transect were: wetted width, three channel depth measurements, % large woody debris and riparian shading. On-site habitat assessments were conducted using the rapid assessment protocol developed for the EPA by Barbour et al. (1999) with modifications for the BLM by the National Aquatic Assessment Team (scores 0-24) (BLM 2008). The process for determining Proper Functioning Condition followed Pritchard et al. (1993). Basic water parameters (temperature, pH, conductivity) were recorded prior to biological sampling using a Horiba H-10 water monitor, calibrated for the higher conductivity range. The Livestock Use Index (“Cowpie” CPI) was assessed by walking a randomly chosen 75m reach on both sides of the stream channel in the riparian area within the assessment area and counting all the old and new cowpies (high CPI equals high cow usage). The goal of these evaluations is to characterize local reach geomorphology, riparian and

in-stream habitat, and characteristics that influence aquatic community integrity. The sites ranking higher using these protocols are determined to have higher quality local-scale habitat. Habitat assessments were performed during the same visit as the biological sampling.

Macroinvertebrate Communities

Macroinvertebrate communities were sampled qualitatively from the 10 transects within the 300m assessment reach using the EMAP Reach-Wide protocol (Lazorchak et al. 1998). Sampling started at the downstream transect (A) or #10 in the BLM protocol, and proceeded upstream alternating sampling with the 500-micron D-frame net to the left, right or center of the stream channel, so a random sampling of all habitats is achieved (Figure 2). Ten multi-habitat kicks were composited from the net into a 20 liter bucket.



Figure 2. Macroinvertebrate EMAP Sampling procedure .

All organisms and organic matter in the bucket were elutriated from the inorganic portion and washed onto a 500-micron sieve. The inorganic portion was washed and examined until no further organics or organisms were present and discarded. The organic portion on the sieve was transferred to one or two 1 liter Nalgene bottles (unless field sub-sampling was needed), labeled and preserved in 95% ethanol and brought to the MTNHP lab in Helena for processing (sorting, identification and data analysis) following

protocols outlined by the BLM (2008a) and MTDEQ (2006). Macroinvertebrates were identified to the lowest taxonomic level (MTDEQ 2006), counted, imported into EDAS (Jessup 2006), and biological metrics were calculated from the data using the Montana Department of Environmental Quality's newest multimetric macroinvertebrate (MMI) protocols (Jessup et al. 2005, Feldman 2006). Metric results were scored using the MTDEQ bioassessment criteria and each sample categorized as nonimpaired or impaired according to threshold values (Table 2). The macroinvertebrate MMI score is based upon a series of metrics that measure attributes of benthic

Table 2. Impairment determinations from the MTDEQ MMI and O/E (RIVPACS) models

Ecoregion	RIVPACS	MMI	Impairment Determination
Mountain	≥ 0.8 or ≤ 1.2	≥ 63	Not impaired
	< 0.8 or > 1.2	< 63	Impaired
Low Valley	≥ 0.8 or ≤ 1.2	≥ 48	Not impaired
	< 0.8 or > 1.2	< 48	Impaired
Eastern Plains	≥ 0.8 or ≤ 1.2	≥ 37	Not impaired
	< 0.8 or > 1.2	< 37	Impaired

macroinvertebrate communities that are sensitive to condition changes in the stream (in the form of pollution or pollutants). The index score represents the condition of the macroinvertebrate community at the time the sample was collected within that past year. If the index score is below the impairment threshold, the individual metrics can be used to provide insight as to why the communities are different from the reference condition (Barbour et. al 1999, Jessup et. al. 2005).

The impairment threshold set by MTDEQ is **37** for the Eastern Plains Stream Index, thus any scores above this threshold are considered unimpaired.

Fish and Amphibian Surveys

Fish sampling within the 300m stream channel assessment reach was conducted with 6 and 9 meter straight seines in 25-30m increments seining in a downstream direction toward the block seine (protocols outlined in Bramblett et al. 2005). Fish captured in a blocked section were transferred to holding buckets until the reach was completed, unless the reach was broken up by riffles, impassable or dry sections; in this case, fish were worked up and released within the section of capture. Fish holding in the buckets were identified to species (Holton and Johnson 2003), enumerated, examined for external anomalies (e.g. deformities, eroded fins, lesions, and



Figure 3. Brassy minnows and lake chubs in seine-Otter 22.

tumors), and then released. At least 10% of the individuals of a species were measured for total length in millimeters (TL mm) to obtain size structure data. Young-of-the-year fish less than 20 mm (TL) were noted on the field sheet (not included in the totals) and released. Baited minnow traps were deployed and fished overnight (12

hour minimum) during the spring visits when seining conditions were toughest. Voucher specimens were only taken in the case of uncertain field identifications, and were preserved in 10% buffered formalin. These will be deposited with the Montana State University Collections. Adult amphibians or reptiles seen while seining or walking the designated stream reach were counted and recorded even if they were not captured in the seine.

Analysis of the sampled fish communities used Integrated Biotic Indices (IBI) designed for wadable prairie streams (Bramblett et. al 2005) and derived Observed/Expected (O/E) Fish Models (Stagliano 2011) to detect impairment in the biological integrity of the sites. The expected number of native fish species for a D005 classified reference stream is 2.5-3.75, while the number of fish for a C005 stream is 5.5-8.5 depending on watershed area, dividing the observed number of native fish species at a site by the expected number derives a percentage compared to reference

condition (>0.8 or 80% = unimpaired) (Table 1). The IBI originally proposed by Karr (1981) involved the calculation of a series of 12 metrics evaluating different attributes of the fish community (i.e. species richness, tolerance to pollutants, trophic status) (Table 3). The 10 metrics used for the prairie streams were adjusted for watershed area to calculate an overall score between 0 and 100. Bramblett et al. (2005) did not propose threshold criteria for good, fair, and poor biological integrity for these scores. Therefore, we applied percentiles above the null criteria (no fish present score) at >30% indicates good to excellent biological integrity, 10-30% fair/good biological integrity, 0-10% indicated poor to fair biological integrity and scores below the null are indicative of poor biological integrity or severely impaired.

Table 3. Fish metrics and classification of fishes captured during the Otter Creek Study (2011).

Species	Scientific Name	Trophic *	Feeding Habitat†	Repro Guild‡	General Tolerance **	Origin ††	Total Length 3 years
Catostomidae							
White sucker	<i>Catostomus commersoni</i>	OM	BE	LO	TOL	N	229
Cyprinidae							
Common Carp	<i>Cyprinus carpio</i>	OM	BE		TOL	I	381
Brassy minnow	<i>Hybognathus hankinsoni</i>	HB	BE		MOD	N	81
Fathead Minnow	<i>Pimephales promelas</i>	OM	GE	TOL§	TOL	N	76
Lake Chub	<i>Couesius plumbeus</i>	OM	GE		MOD	N	140
Sand Shiner	<i>Notropis stramineus</i>	OM	GE	LO	MOD	N	61
Centrarchidae							
Green Sunfish	<i>Lepomis cyanellus</i>	IC	GE	TOL§	TOL	I	102
Pumpkinseed	<i>Lepomis gibbosus</i>	IC	GE	TOL§	MOD	I	89
Ictaluridae							
Black Bullhead	<i>Ameiurus melas</i>	IC	BE	TOL§	TOL	I	152

*HB = herbivore (> 90% plants or detritus); IC = invertivore/carnivore (>25% both invertebrates and vertebrates); IN = invertivore; OM = omnivore(25-90% plants or detritus)

† BE = benthic; GE = generalist; WC = water column; Brown (1971); Scott and Crossman (1973); Becker (1983)

‡ LO=Litho-obligate Reproductive Guild; Scott and Crossman (1973); Pflieger (1997); Barbour et al. (1999)

§ Tolerant reproductive strategists are not litho-obligates, use parental care at spawning site; Scott and Crossman (1973); Pflieger (1997)

** INT = intolerant; MOD = moderately tolerant; TOL = tolerant; Barbour et al. (1999);

†† N = native; I – introduced; Brown (1971); Holton and Johnson (2003)

Results

We evaluated seven stream reaches in the study area: four Otter Creek mainstem sites were classified as Great Plains Perennial Prairie Stream types (C005), and the three tributaries were the Great Plains Intermittent Prairie Stream systems (D005, E005) (Stagliano 2005). Proper

classification is important when determining biological integrity (Hawkins and Norris 2000) and expected species richness. The Intermittent Prairie Stream (E005) in Montana is naturally fishless about 80% of the time; therefore, absence of fish, in itself, should not be viewed as a biological impairment (Threemile and Tenmile Creek). Likewise, stream reaches of Otter Creek have become dry in previous years (Stagliano, personal observation 2005-2008), thus placing this stream section within the D005 classification. We have identified and characterized reference condition indicator assemblages for these ecosystem types previously (Stagliano 2005), and used these here to compare to our site-specific observed species.

Habitat Evaluations. Of the seven sampling reaches evaluated within the study area, we found three in Proper Functioning Condition (PFC) with a stable trend and four were Functional at Risk (FAR) (Table 7). Reasons that sites ranked FAR were due to structural habitat alteration by cattle with associated high CPI values (Home Creek {Otter_1A}, Threemile Creek {Otter_3m} and Otter #16-fall) (Figure 4) or anthropogenic stream manipulation (Otter Creek JT-Trussler and Otter Creek

Figure 4. Structurally cattle damaged reach-Home Creek



#16). Highest site integrity scores using both the BLM Habitat and PFC Assessment methods were recorded at the Tenmile Creek (Otter_23) and Otter Creek #22 (control-Denson reach) (Table 7). Point conductivity measurements recorded at all Otter Creek mainstem sites and tributaries across all seasons were above the threshold for impairment levels ($>500\mu\text{s}$, DEQ 2006), and Home Creek site had visible signs of natural gas seepage from the sediments.


Macroinvertebrate Communities: Overall, 104 unique macroinvertebrate taxa were reported from the 15 macroinvertebrate assessment samples (Appendix B). One known Montana species of concern (SOC), the mayfly, *Caenis youngi* was collected in fair numbers at the control site, Otter Creek #22. Stoneflies (P) were not present at any sites, so the EPT taxa per site usually consisted of two species of tolerant mayflies (E), and one or two species of caddisfly (T); the highest EPT richness at any site was six species at the Otter JT site spring (Table 4, Appendix B). Average

macroinvertebrate taxa richness per site was 29.5 and the highest taxa richness was 41 taxa reported at the Otter Creek JT site. Using the MTDEQ multimetric index (MMI), four of the five sites (12 of 15 samples) were ranked non-impaired (good to excellent biological integrity), while all three samples from Tenmile Creek were ranked marginally impaired (Figure 5).

Table 4. Macroinvertebrate sample characteristics and various metrics used for the DEQ MMI.


Site ID	CollDate	Total Ind. ID'ed	Total Sample	Total Taxa	Plains MMI Index	EPT Taxa	% EPT	HBI	% Non- Insect
Otter_23	5/16/2011	605	4,840	15	36.8	1	0.3	7.7	6.9
Otter_23s	7/17/2011	515	2,060	12	35.2	0	0.0	7.3	15.2
Otter_23f	10/15/2011	617	2,468	13	35.6	0	0.0	7.1	15.9
OTTER_22	5/17/2011	628	5,024	24	46.3	3	2.7	7.4	56.2
OTTER_22s	7/14/2011	879	10,548	35	54.3	3	36.2	7.9	40.7
OTTER_22f	10/14/2011	725	5,800	35	65.9	3	22.3	7.7	43.5
OTTER_16	5/17/2011	564	4,512	34	44.4	3	3.2	7.5	55.3
OTTER_16s	7/16/2011	686	8,232	35	51.2	2	25.4	7.7	43.0
OTTER_16f	10/15/2011	603	4,824	35	46.2	4	22.9	7.2	50.3
OTTER_JT	5/17/2011	610	2,440	34	64.6	6	17.4	7.4	30.2
OTTER_JTs	7/18/2011	605	3,630	28	65.8	5	18.2	7.5	37.7
OTTER_JTf	10/15/2011	606	2,424	41	43.4	4	7.6	6.6	10.9
OTTER_1A	5/16/2011	639	5,112	35	57.4	2	2.5	6.8	17.8
OTTER_1As	7/17/2011	626	5,008	32	43.2	1	0.2	7.2	11.8
OTTER_1Af	10/14/2011	617	4,936	35	37.9	1	0.3	7.0	21.7

Figure 5. DEQ MMI scores across sites and seasons. Line is the impairment threshold at 37.



Stream sites that maintained flowing, connected water scored higher with the MMI than sites with interrupted pool areas. Overall, mainstem sites evaluated in the Otter Creek study received significantly higher macroinvertebrate MMI scores than those in the tributaries (T-test, $p < 0.01$). MMI's did not significantly differ on Otter Creek mainstem Pre-Impact Control, Impact or Downstream Sites (T-test, $p > 0.05$), despite the fish communities reflecting a decrease in biotic integrity. No site had consistently high macroinvertebrate scores across all seasons.

Figure 6. Average MMI scores by site type. (a) = no baseline differences between treatments.



Fish Communities. Overall, nine fish species (five native/four introduced) were identified from 37,679 individuals collected from 15 site visits (Table 5). One potential species of concern (PSOC), the brassy minnow, was collected at five of six sites and in very high numbers at the control site, Otter 22. Dominance by brassy minnows and lake chubs at the upstream sites shifts to a tolerant fathead minnow and sand shiner dominated community at the downstream site, Otter JT (Table 5). Otter Creek site 16 was heavily infested by introduced species during the summer and fall visits representing 41% and 93% of the fish sampled, respectively. Average number of fish species collected at Otter Creek mainstem sites across all seasons was 6.5 (± 0.8 SE), while the tributary sites averaged 1.75 species (Table 5). All fish presence sites also reported at least one species of amphibian (Table 6). Lake chubs had the highest site occupancy rate at 93% (14 of 15 visits) followed by fathead and brassy minnows at 80% (12 of 15 visits), but fathead minnows contributed to the highest proportion of individuals (56%, see Table 5), followed by lake


chubs at 17%. The most diverse site in the study was Otter Creek JT site with nine species, while the most intact sites were Otter Creek Site #22 at Denson's (four native spp.) and the Tenmile Creek spring survey (two native spp.).

Table 5. Fish abundance, IBI's and O/E results for the 6 sites reporting fish. ns =not seined during visit (dry) or us =unseivable conditions. Underlined values are fish communities ranked biologically unimpaired.

	OTTER_23			OTTER_22			OTTER_16			OTTER_2			OTTER_JT			OTTER_1A			
	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	Total
Black Bullhead*	0	ns	ns	0	5	3	0	0	186	us	6	7	13	3	5	0	0	0	228
Brassy Minnow	25	ns	ns	81	2400	588	104	549	51	us	154	364	143	30	65	0	0	0	4554
Common Carp*	0	ns	ns	0	0	0	4	432	1314	us	14	49	20	213	138	0	0	0	2184
Fathead Minnow	0	ns	ns	0	0	3	40	3	12	us	56	35	2008	2995	17810	10	5	1	22978
Lake Chub	12	ns	ns	153	2785	1059	16	84	9	us	105	84	5	8	103	22	12	0	4457
Green Sunfish*	0	ns	ns	0	0	3	0	0	6	us	0	0	3	3	3	0	0	0	18
Pumpkinseed*	0	ns	ns	0	0	3	0	3	30	us	7	0	0	0	10	0	0	0	53
Sand Shiner	0	ns	ns	0	0	0	0	0	0	us	22	0	38	25	2755	0	0	0	2840
White Sucker	0	ns	ns	6	140	15	20	1	45	us	8	7	28	108	10	0	0	0	388
Total # species	2	0	0	3	4	7	5	6	8	0	8	6	8	8	9	2	2	1	9
Native Species	2	0	0	3	3	4	4	4	4	0	5	4	5	5	5	2	2	1	5
Total Individuals	37	0	0	240	5330	1674	184	1072	1653	0	372	546	2257	3384	20899	32	17	1	37,698
IBI	<u>75.6</u>			<u>66.1</u>	<u>67.3</u>	<u>69.2</u>	<u>62.3</u>	50.7	44.4		<u>62.7</u>	<u>63.1</u>	44.9	43.8	48.9	<u>70.8</u>	<u>70.8</u>	46.5	
O/E	<u>0.82</u>			<u>0.80</u>	<u>0.80</u>	<u>1.07</u>	0.73	0.73	0.73		<u>0.91</u>	0.73	0.59	0.59	0.59	<u>0.82</u>	<u>0.82</u>	0.41	
O/E %	<u>81.6</u>			<u>80.0</u>	<u>80.0</u>	<u>106.7</u>	72.7	72.7	72.7		<u>90.9</u>	72.7	58.8	58.8	58.8	<u>81.6</u>	<u>81.6</u>	40.8	

Using the Prairie Fish IBI, 9 of the 15 fish sites were ranked non-impaired (good biological integrity), two were slightly impaired (moderate integrity), three were moderately impaired (poor biotic integrity) and the Home Creek fall survey was ranked severely impaired (Figure 7). The O/E models tracked the IBI scores quite well at most sites (13 of 15), except two where the O/E ranked the site as slightly impaired and the IBI ranked the sites as non-impaired (Figure 8).

Figure 7. Fish IBI scores across sites and seasons. Line is the null IBI threshold (fish absent).



Fish IBI's decreased going downstream on Otter Creek, and the Pre-Impact Control Site scored significantly higher than either the Impact or Downstream sites (T-test, $p = 0.05$) (Figure 9). Within the Great Plains Intermittent Prairie Streams, the species expected to occur most often at reference condition sites are the fathead minnow and brook stickleback, but the upper Tongue and Powder River Basins are slightly out of the range of this latter species; therefore, we did not collect them or expect them to occur at these sites and revised the Observed /Expected (O/E) Fish Model predictors accordingly. The O/E scores agreed with the IBI scores in most cases,

except for the Otter Creek stream reaches that could be classified as Perennial vs. Intermittent Prairie Streams (C005) (Table 5).

Figure 8. Fish IBI compared to O/E model (%) scores across sites and seasons.





Figure 9. Average fish IBI scores by site type. (a) = no baseline differences between treatments.



Amphibian and Reptile Incidentals. Overall, eight herpetofauna species were observed, collected in dipnets/seines or incidentally recorded in conjunction with the fish, habitat and macroinvertebrate surveys. Otter Creek #22 and #16 were the most diverse sites reporting five herpetofuana species cumulatively across all seasons. We reported four amphibian species, of which Woodhouse's Toad (*Bufo woodhousii*) had the highest site occupancy, occurring at six of seven sites, followed by the Northern Leopard Frog (*Rana pipiens*) and Tiger Salamander (*Ambystoma tigrinum*) (Figure 10) recorded at five and four sites, respectively (Table 6). Boreal Chorus Frog (*Pseudacris maculata*) was detected vocally calling at three sites during the spring visits only and one incidental sighting during a summer visit. We also recorded four reptile species (in order of site occurrence): Western Rattlesnake (*Crotalus viridis*), Painted Turtle (*Chrysemys picta*) (Figure 10), Terrestrial Garter Snake (*Thamnophis elegans*) and Snapping Turtle (*Chelydra serpentina*) (SOC).

Table 6. Vertebrates (Species Code) recorded during the Otter Creek Surveys. Frequency of Occurrence (FO) was calculated from the # of site visits detected / # of visits capable for detection: Herps (n=21) and Fish (n=15). * = Introduced Species

Herpetofauna	Visits Det	FO
Woodhouse's Toad (BUWO)	9	0.43
Northern Leopard Frog (RAPI)	7	0.33
Tiger Salamander (AMTI)	5	0.24
Boreal Chorus Frog (PSMA)	4	0.19
Painted Turtle (CHPI)	4	0.19
Western Rattlesnake (CRVI)	4	0.19
Snapping Turtle (CHSE)	2	0.10
Terrestrial Gartersnake (THEL)	2	0.10
Fish		
Lake Chub (LACH)	14	0.93
Fathead Minnow (FAMI)	12	0.80
Brassy Minnow (BRMI)	12	0.80
White Sucker (WHSU)	10	0.67
Black Bullhead* (BLBU)	8	0.53
Common Carp* (CARP)	8	0.53
Green Sunfish* (GRSU)	5	0.33
Pumpkinseed* (PUMP)	5	0.33
Sand Shiner (SASH)	4	0.27

Figure 10. Tiger Salamander (left) from Home (Otter_1A) and Painted Turtle from OtterJT (right).



Conclusions

The aquatic community sampling sites chosen for pre-impact baseline data were representative of the range of stream classes found in the Otter Coal Tracts project area: Ephemeral, Intermittent and Perennial Prairie Streams. Despite this being an unusually high water year for the region, stream communities that we sampled across the seasons encompassed the range of expected species to occur in these stream types, and fish and macroinvertebrate community assessment scores were similar to biological assessments performed in previous years (Table 7).

Table 7. Otter Creek Temporal Habitat Scores, Fish IBIs and MMIs. Metric Score Ranks (+): unimpaired, (-): impaired, (=): marginal, (na)= not applicable, taxa group not collected. Spatial relationship in river miles (rm) to them adjacent Coal Tract Study Site

Site Code	Site Name	Spatial Relation	Date	BLM PFC	BLM Hab Score	Fish IBI	MT MMI	Metric Score Ranks
ATG_203	Otter Creek at Camp Creek Road	~16 rm ups	7/11/2010	PFC	20	50	58.4	+, +, +
ATG_235	South Fork Taylor Creek	~13 rm ups	7/11/2010	PFC	22	71	73.5	+, +, +
ATG_202	Otter Creek at Taylor Creek Road	~12 rm ups	7/11/2010	PFC	19	60	36.3	+, +, -
ATG_CCC	Otter Creek at CCC Camp USFS	~3 rm ups	7/11/2010	PFC	19	58	37.9	+, +, =
ATG_201	Otter Creek at Tenmile Cr Road	100m ups	7/11/2010	PFC	19	52	44.6	+, =, +
OTTER_23	Tenmile Creek @ Denson Ranch	0 rm	5/10/2011	FAR	16	75	36.8	-, +, -
OTTER_23	Tenmile Creek @ Denson Ranch	0 rm	7/18/2011	FAR	17	na	35.2	-, na, -
OTTER_23	Tenmile Creek @ Denson Ranch	0 rm	10/10/2011	FAR	17	na	35.6	-, na, -
OTTER_22	Otter Creek at Tenmile Cr Road	0 rm	5/10/2011	PFC	22	66.1	46.3	+, +, +
OTTER_22	Otter Creek at Tenmile Cr Road	0 rm	7/18/2011	PFC	22	67.3	54.3	+, +, +
OTTER_22	Otter Creek at Tenmile Cr Road	0 rm	10/10/2011	PFC	22	69.2	65.9	+, +, +
OTTER_16	Otter Creek at Ranch Road x	0 rm	5/19/2011	PFC	19	62.3	44.4	+, +, +
OTTER_16	Otter Creek at Ranch Road x	0 rm	7/16/2011	PFC	19	50.7	51.2	+, -, +
OTTER_16	Otter Creek at Ranch Road x	0 rm	10/15/2011	FAR	18	44.4	46.2	=, -, +
OTTER_2	Otter Creek at Ranch Road x	0 rm	7/16/2011	FAR	17	62.7	na	=, +, na
OTTER_2	Otter Creek at Ranch Road x	0 rm	10/15/2011	FAR	17	63.1	na	=, +, na
YL_S70REM	Otter Creek at 3mile creek	400m dns	7/18/2000	na	na	na	59.7	na, na, +
OTTER_JT	Otter Creek J Trusler site	0 rm	5/17/2011	PFC	19	44.9	64.6	+, -, +
OTTER_JTs	Otter Creek J Trusler site	0 rm	7/18/2011	PFC	19	43.8	65.8	+, -, +
OTTER_JTf	Otter Creek J Trusler site	0 rm	10/15/2011	PFC	19	48.9	43.4	+, -, +
YL_S0070dn	Otter Creek below Trusler site	300m dns	6/2/2003	na	na	na	56.2	na, na, +
OTTER_1A	Home Creek	0 rm	5/16/2011	FAR	11	70.8	57.4	-, +, +
OTTER_1A	Home Creek	0 rm	7/17/2011	FAR	12	70.8	43.2	-, +, +
OTTER_1A	Home Creek	0 rm	10/15/2011	FAR	13	46.5	37.9	-, -, =

Biotic integrity of the upstream control reaches of Otter Creek was initially higher during this baseline study. One extraordinary finding was the high density and large biomass of fish inhabiting the stream reach below Trusler's Ranch road crossing (20,000 fish in the 300 m reach

fall survey, 15 fish/m²); this is largely an artifact of the impassible culvert located here and the fish “stacking up” downstream of this barrier. Density dependent fish anomalies (lesions and parasites, i.e., yellow grub and anchor worm) are prevalent at this site, indicating the fish are likely experiencing stress from overcrowding. Benthic macroinvertebrate densities at this site are low, but, more diverse than other sites, indicating that the fish may be consuming large portions of the available insect production having a top-down community effect. Outside of coal extraction, threats to this stream system, which can be managed, include grazing and livestock use around the riparian areas. Moderate occurrences in these basins can have strong local effects resulting in sedimentation, stream widening at cattle crossings and loss of functional channel hydrology. Introductions of game (green sunfish, bullheads or pumpkinseeds) or forage fish in stock ponds anywhere in the watershed can pose potential problems for native fish as these introduced fish become permanent residents, outcompete or prey upon the native fish and contribute to overall community degradation. Diverse aquatic communities with high biological integrity are usually correlated with good riparian condition and habitat quality (Allen et al. 1997). Thus, effective riparian zone management of cattle grazing would contribute to intact vegetation buffers and less sediment in the aquatic environment (George et al. 2002). In this study, macroinvertebrate communities ranked few sites as impaired, even those with an obvious impaired riparian condition, but the effectiveness of macroinvertebrates in assessing prairie stream impairment is still under debate in Montana. Community results from the habitat, fish and macroinvertebrate surveys combined to rank the following sites from highest biological integrity to lowest within their aquatic ecological classification codes:

Northwestern Great Plains Perennial Prairie Stream (AES code C005)-1) Otter Creek #22, 2)
Otter Creek #2, 3) Otter Creek #16, 4) Otter Creek –J Trusler site

Site Photos

Otter Creek #22 (control) during spring visit: pools (left) and riffle (right)



Otter Creek #2 (impact): fall visit looking downstream to block net (left) and upstream (right).



Site Photos

Otter Creek #16 (impact) spring visit: pool (left) and shallow stream crossing (right).



Otter Creek #16 (impact) during summer visit: downstream pool (left) and drill trucks using the stream crossing (right).



Site Photos

Otter Creek JT (downstream): summer visit looking downstream (left) and upstream (right)



Northwestern Great Plains Intermittent Prairie Stream-(AES code D005)-1) Home Creek 1A, 2)
Tenmile Creek,

Home Creek (Otter 1A) during the Spring (left) and Summer (right) Visits



Site Photos

Tenmile Creek (Otter_23) in the Spring (left) and Summer (right)



Great Plains Intermittent Fishless Prairie Stream (AES code E005)-1) Threemile Creek

Site Photos: Threemile Creek (Otter 3m) during the spring (left) and summer (right) Visits



Literature Cited

- Allan, J. D., D. L. Erickson and J. Fay. 1997. The Influence of Catchment Land Use on Stream Integrity Across Multiple Spatial Scales. *Freshwater Biology* 37:149-162.
- Barbour, M., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadable Rivers: Periphyton, Benthic Macroinvertebrates and Fish, Second Edition. EPA 841-B-99-002. United States Environmental Protection Agency; Office of Water: Washington, D.C.
- BLM 2008. Standard Methods for Laboratory Sample Sorting Procedures of Macroinvertebrate Samples. <http://usu.edu/buglab/SampleProcessing/labProcedures.cfm> Bureau of Land Management/USU National Aquatic Monitoring Center, Logan, Utah.
- BLM 2008b. Standard Methods for Field Surveys of Macroinvertebrate Samples. <http://www.usu.edu/buglab/Monitoring/fieldForms.cfm> Bureau of Land Management/USU National Aquatic Monitoring Center, Logan, Utah.
- Bramblett, R. G., T. R. Johnson, A. V. Zale, A. V., and D. Heggem. 2005. Development and Evaluation of a Fish Assemblage Index of Biotic Integrity for Northwestern Great Plains. *Transactions of the American Fisheries Society* 134:624–640, 2005.
- Bramblett, R. G., and K. D. Fausch. 1991. Variable Fish Communities and the Index of Biotic Integrity in a Western Great Plains River. *Transactions of the American Fisheries Society* 120:752–769.
- Dodds, W. K., K. Gido, M. R. Whiles, K. M. Fritz, and W. J. Matthews. 2004. Life on the Edge: The Ecology of Great Plains Prairie Streams. *BioScience* 54: 205–216.
- DOI Bureau of Land Management 2002. Environmental Assessment. <http://dnrc.mt.gov/Trust/MMB/OtterCreek/2General/TransferDocuments/BLMTransferEA.pdf>
- Feldman, D. 2006. Interpretation of New Macroinvertebrate Models by WQPB. Draft Report. Montana Department of Environmental Quality, Planning Prevention and Assistance Division, Water Quality Planning Bureau, Water Quality Standards Section. 1520 E. 6th Avenue, Helena, MT 59620. 14 pp.
- George, M.R., R.E. Larsen, N.K. McDougald, K.W. Tate, J.D. Gerlach, Jr., and K.O. Fulgham. 2002. Influence of Grazing on Channel Morphology of Intermittent Streams. *J. Range Management*. 55:551-557.
- Hawkins, C. P. and R. H. Norris. 2000. Performance of different landscape classifications for aquatic bioassessments: introduction to the series. *Journal of the North American Benthological Society*. 19:3 (367-369).

- Holton, G. D., and H. E. Johnson. 2003. A Field Guide to Montana Fishes, 3rd edition. Montana Fish, Wildlife, and Parks, Helena.
- Jessup, B., J. Stribling; and C. Hawkins. 2005. Biological Indicators of Stream Condition in Montana Using Macroinvertebrates. Tetra Tech, Inc. November 2005 (draft).
- Jessup, B. 2006. Ecological Data Application System (EDAS) Version MT 3.3.2k A User's Guide. Tetra Tech, Inc.
- Karr, J. R. 1981. Assessment of Biotic Integrity Using Fish Communities. Fisheries 6(6):21–27.
- Lazorchak, J.M., Klemm, D.J., and D.V. Peck (editors). 1998. Environmental Monitoring and Assessment Program - Surface Waters: Field Operations and Methods for Measuring the Ecological Condition of Wadeable Streams. EPA/620/R-94/004F. U.S. Environmental Protection Agency, Washington, D.C.
- Montana Department of Environmental Quality (DEQ). 2006. Sample Collection, Sorting, and Taxonomic Identification of Benthic Macroinvertebrates. Water Quality Planning Bureau. Standard Operation Procedure (WQPBWQM-009).
- Montana Department of Environmental Quality (DEQ). 2006. ADMINISTRATIVE RULES OF MONTANA 17.30.670. Chapter 30 Surface Water Quality Standards and Procedures. www.deq.mt.gov/dir/Legal/Chapters/Ch30-06.pdf
- Peterson, D.A., Clark, M.L., Foster, Katharine, Wright, P.R., and Boughton, G.K., 2010, Assessment of Ecological Conditions and Potential Effects of Water Produced From Coalbed Natural Gas Development on Biological Communities in Streams of the Powder River Structural Basin, Wyoming and Montana, 2005–08: USGS Scientific Investigations Report 2010–5124, 84 p. <http://pubs.usgs.gov/sir/2010/5124/pdf/SIR10-5124.pdf>
- Pritchard, D., F. Barret, H. Berg, W. Hagenbuck, R. Krapf, R. Leinard, S. Leonard, M. Manning, C. Noble, Tippy, D. 1993. Riparian Area Management: A User Guide to Assessing Proper Functioning Condition. Technical Reference 1737-9. USDI Bureau of Land Management Service Center. Denver, Colorado. USA. 109 pp.
- Stagliano, David, M. 2011. Baseline Assessments for Fish, Macroinvertebrates, and Herpetofauna in the Headwaters of Otter and Hanging Woman Creeks within the Tongue Powder CBNG Area. Report to the Bureau of Land Management. Montana Natural Heritage Program, Helena, Montana. 17 pp. [http://mtnhp.org/reports/CBNG TonguePowder.pdf](http://mtnhp.org/reports/CBNG_TonguePowder.pdf)
- Stagliano, David, M. 2005. Aquatic Community Classification and Ecosystem Diversity in Montana's Missouri River Watershed. Report to the Bureau of Land Management. Montana Natural Heritage Program, Helena, Montana. 65 pp. plus appendices. <http://www.mtnhp.org/reports.asp#Ecology>

Appendix A. Raw data and IBI metric calculation from fish data collected from Otter Creek Coal Tract Sites									
	Spring 2011								
	Otter23		Otter22		Otter16		OtterJT		Otter1A
Black Bullhead	0.0		0.0		0.0		12.5		0.0
Brassy Minnow	25.0		81.0		100.0		142.5		0.0
Common Carp	0.0		0.0		4.0		20.0		0.0
Fathead Minnow	0.0		0.0		16.0		2007.5		10.0
Lake Chub	12.0		153.0		40.0		5.0		20.0
Green Sunfish	0.0		0.0		0.0		2.5		0.0
Pumpkinseed Sunfish	0.0		0.0		0.0		0.0		0.0
Sand Shiner	0.0		0.0		0.0		37.5		0.0
White Sucker	0.0		6.0		20.0		27.5		0.0
Total # species	2.0		3.0		5.0		8.0		2.0
Native Species	2.0		3.0		4.0		5.0		2.0
Native Families	1.0		2.0		2.0		2.0		1.0
Total Individuals	37.0		240.0		180.0		2255.0		30.0
# Minnow Species Thrive	2.0		2.0		3.0		4.0		2.0
Proportion of tolerant individuals	0.00		2.50		22.22		91.80		33.33
# Sucker + Catfish Species	0.0		1.0		1.0		2.0		0.0
% Insectivorous Minnows	32.4		63.8		22.2		1.9		66.7
# Benthic Invertivore Species	0.0		0.0		0.0		1.0		0.0
% Lithophilic Spawners	0.0		2.5		11.1		2.9		0.0
% Parental Care	0.00		0.00		8.89		89.58		33.33
% Native to Montana	100.0		100.0		97.8		98.4		100.0
# Long Lived Species	2.0		1.0		2.0		4.0		1.0
	Otter23		Otter22		Otter16		OtterJT		Otter1A
Metrics	Adjust Value	Score	Adjust Value	Score	Adjust Value	Score	Adjust Value	Score	Adjust Value
Number of Native Fish Species to Montana	13.6	75.8	9.3	51.8	10.1	56.0	10.8	59.9	13.3
Number of Native Fish Families to Montana	3.1	58.0	3.2	58.7	3.1	57.8	3.1	56.8	3.1
Proportion of tolerant individuals	0.0	100.0	2.5	97.4	22.2	76.5	91.8	2.9	33.3
Number of Sucker and Catfish Species	7.1	77.1	4.9	53.3	4.8	51.7	5.6	60.7	6.9
Proportion out of the Total Number of Fish That Were Insect eating Minnows	32.4	44.5	63.8	87.6	22.2	30.5	1.9	2.6	66.7
Total Number of Species That Prefer to Eat Insects That Live on the Stream Bottom	5.5	93.1	3.0	51.3	4.9	83.3	4.8	81.0	5.3
Proportion of the Total Number of Fish That Require Rocks to Lay Eggs	0.0	0.0	2.5	3.0	11.1	13.4	2.9	3.5	0.0
Proportion of the Total Number of Individuals That Do Not Require Rocks, But Have Parental Care of Eggs	0.0	100.0	0.0	100.0	8.9	89.9	89.6	-1.9	33.3
Proportion of the Total Number of Fish Sampled That Were Native to Montana	100.0	100.0	100.0	100.0	97.8	97.8	98.4	98.5	100.0
Number of Long-Lived Native Species	10.5	107.4	5.7	58.1	6.5	66.5	8.3	84.9	9.2
		756.0		661.2		623.5		448.9	
IBI Score		75.60		66.12		62.35		44.89	

Appendix A. Raw data and IBI metric calculation from fish data collected from Otter Creek Coal Tract Sites

Summer 2011												
	Otter23		Otter22		Otter16		Otter2		OtterJT		Otter1A	
Black Bullhead	0.0		5.0		0.0		7.0		2.5		0.0	
Brassy Minnow	0.0		2400.0		549.0		154.0		30.0		0.0	
Common Carp	0.0		0.0		432.0		14.0		212.5		0.0	
Fathead Minnow	0.0		0.0		3.0		56.0		2995.0		10.0	
Lake Chub	0.0		2785.0		84.0		105.0		7.5		20.0	
Green Sunfish	0.0		0.0		0.0		0.0		2.5		0.0	
Pumpkinseed Sunfish	0.0		0.0		3.0		7.0		0.0		0.0	
Sand Shiner	0.0		0.0		0.0		22.0		25.0		0.0	
White Sucker	0.0		140.0		0.0		7.0		107.5		0.0	
Total # species	0.0		4.0		5.0		8.0		8.0		2.0	
Native Species	0.0		3.0		3.0		5.0		5.0		2.0	
Native Families	0.0		2.0		1.0		2.0		2.0		1.0	
Total Individuals	0.0		5330.0		1071.0		372.0		3382.5		30.0	
# Minnow Species Thrive	0.0		2.0		3.0		4.0		4.0		2.0	
Proportion of tolerant individuals	0.00		2.72		40.62		22.58		98.15		33.33	
# Sucker + Catfish Species	0.0		2.0		0.0		2.0		2.0		0.0	
% Insectivorous Minnows	0.0		52.3		7.8		34.1		1.0		66.7	
# Benthic Invertivore Species	0.0		1.0		0.0		1.0		1.0		0.0	
% Lithophilic Spawners	0.0		2.6		0.0		7.8		3.9		0.0	
% Parental Care	0.00		0.09		0.28		16.94		88.62		33.33	
% Native to Montana	0.0		99.9		59.4		92.5		93.6		100.0	
# Long Lived Species	0.0		1.0		2.0		1.0		4.0		1.0	
	Otter23		Otter22		Otter16		Otter2		OtterJT		Otter1A	
Metrics	Adjust Value		Adjust Value		Adjust Value		Adjust Value		Adjust Value		Adjust Value	Score
Number of Native Fish Species to Montana	11.6	64.7	9.3	51.8	9.1	50.4	11.1	61.5	10.8	59.9	13.3	73.7
Number of Native Fish Families to Montana	2.1	39.5	3.2	58.7	2.1	39.4	3.1	57.8	3.1	56.8	3.1	56.7
Proportion of tolerant individuals	0.0	100.0	2.7	97.1	40.6	57.0	22.6	76.1	98.2	-3.9	33.3	64.7
Number of Sucker and Catfish Species	7.1	77.1	5.9	64.2	3.8	40.8	5.8	62.6	5.6	60.7	6.9	74.7
Proportion out of the Total Number of Fish That Were Insect eating Minnows	0.0	0.0	52.3	71.8	7.8	10.8	34.1	46.9	1.0	1.3	66.7	91.6
Total Number of Species That Prefer to Eat Insects That Live on the Stream Bottom	5.5	93.1	4.0	68.3	4.9	83.3	4.9	83.3	4.8	81.0	5.3	90.2
Proportion of the Total Number of Fish That Require Rocks to Lay Eggs	0.0	0.0	2.6	3.2	0.0	0.0	7.8	9.4	3.9	4.7	0.0	0.0
Proportion of the Total Number of Individuals That Do Not Require Rocks, But Have Parental Care of Eggs	0.0	100.0	0.1	99.9	0.3	99.7	16.9	80.7	88.6	-0.8	33.3	62.1
Proportion of the Total Number of Fish Sampled That Were Native to Montana	0.0	0.0	99.9	99.9	59.4	59.4	92.5	92.5	93.6	93.6	100.0	100.0
Number of Long-Lived Native Species	8.5	86.9	5.7	58.1	6.5	66.5	5.5	56.3	8.3	84.9	9.2	94.5
		561.4		673.0		507.4		627.2		438.3		708.4
IBI Score		56.14		67.30		50.74		62.72		43.83		70.84

Appendix A. Raw data and IBI metric calculation from fish data collected from Otter Creek Coal Tract Sites

Fall 2011												
	Otter23		Otter22		Otter16		Otter2		OtterJT		Otter1A	
Black Bullhead	0.0		3.0		186.0		7.0		5.0		0.0	
Brassy Minnow	0.0		588.0		51.0		364.0		65.0		0.0	
Common Carp	0.0		0.0		1314.0		49.0		137.5		0.0	
Fathead Minnow	0.0		3.0		12.0		35.0		17810.0		10.0	
Lake Chub	0.0		1059.0		9.0		84.0		102.5		0.0	
Green Sunfish	0.0		3.0		6.0		0.0		2.5		0.0	
Pumpkinseed Sunfish	0.0		3.0		30.0		0.0		10.0		0.0	
Sand Shiner	0.0		0.0		0.0		0.0		2755.0		0.0	
White Sucker	0.0		15.0		45.0		7.0		10.0		0.0	
Total # species	0.0		7.0		8.0		6.0		9.0		1.0	
Native Species	0.0		4.0		4.0		4.0		5.0		1.0	
Native Families	0.0		2.0		2.0		2.0		2.0		1.0	
Total Individuals	0.0		1674.0		1653.0		546.0		20897.5		10.0	
# Minnow Species Thrive	0.0		3.0		3.0		3.0		4.0		1.0	
Proportion of tolerant individuals	0.00		1.43		94.56		17.95		85.97		100.00	
# Sucker + Catfish Species	0.0		2.0		2.0		2.0		2.0		0.0	
% Insectivorous Minnows	0.0		63.3		0.5		15.4		13.7		0.0	
# Benthic Invertivore Species	0.0		1.0		1.0		1.0		1.0		0.0	
% Lithophilic Spawners	0.0		0.9		2.7		1.3		13.2		0.0	
% Parental Care	0.00		0.36		11.98		7.69		85.25		100.00	
% Native to Montana	0.0		99.5		7.1		89.7		99.3		100.0	
# Long Lived Species	0.0		4.0		4.0		4.0		4.0		1.0	
	Otter23		Otter22		Otter16		Otter2		OtterJT		Otter1A	
Metrics	Adjust Value		Adjust Value		Adjust Value		Adjust Value		Adjust Value		Adjust Value	Score
Number of Native Fish Species to Montana	11.6	64.7	10.3	57.4	10.1	56.0	10.1	56.0	10.8	59.9	12.3	68.2
Number of Native Fish Families to Montana	2.1	39.5	3.2	58.7	3.1	57.8	3.1	57.8	3.1	56.8	3.1	56.7
Proportion of tolerant individuals	0.0	100.0	1.4	98.5	94.6	-0.1	17.9	81.0	86.0	9.0	100.0	-5.8
Number of Sucker and Catfish Species	7.1	77.1	5.9	64.2	5.8	62.6	5.8	62.6	5.6	60.7	6.9	74.7
Proportion out of the Total Number of Fish That Were Insect eating Minnows	0.0	0.0	63.3	86.9	0.5	0.7	15.4	21.1	13.7	18.8	0.0	0.0
Total Number of Species That Prefer to Eat Insects That Live on the Stream Bottom	5.5	93.1	4.0	68.3	4.9	83.3	4.9	83.3	4.8	81.0	5.3	90.2
Proportion of the Total Number of Fish That Require Rocks to Lay Eggs	0.0	0.0	0.9	1.1	2.7	3.3	1.3	1.5	13.2	16.0	0.0	0.0
Proportion of the Total Number of Individuals That Do Not Require Rocks, But Have Parental Care of Eggs	0.0	100.0	0.4	99.6	12.0	86.4	7.7	91.3	85.2	3.0	100.0	-13.7
Proportion of the Total Number of Fish Sampled That Were Native to Montana	0.0	0.0	99.5	99.5	7.1	7.1	89.7	89.8	99.3	99.3	100.0	100.0
Number of Long-Lived Native Species	8.5	86.9	5.7	58.1	8.5	87.0	8.5	87.0	8.3	84.9	9.2	94.5
		561.4		692.2		444.2		631.4		489.4		464.9
IBI Score		56.14		69.22		44.42		63.14		48.94		46.49

Appendix B

Macroinvertebrate taxa lists, abundance and metrics for the Otter
Creek collection sites

Montana Bioassessment Report

Waterbody Name: Otter Creek Site 16 for the Coal Tracts Study **Benthic Sample ID:** 17954
Station ID: OTTER_16 **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTTR16M-MAC-R
Site Classification: **Collection Date:** 05/19/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 564

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Haliphus sp. 2	2			
Amphipoda	Crangonyx	Crangonyx	2		CG	"SW/50%, SP/50%"
Amphipoda	Hyaella	Hyaella	150	8	CG	SW/SP
Basommatop	Lymnaeidae	Pseudosuccinea columella	2	6	CG	CN
Basommatop	Lymnaeidae	Stagnicola caperata	10	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	110	8	CG	CN
Basommatop	Planorbidae	Gyraulus deflectus	2	6	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	24	6	CG	CN
Coleoptera	Berosus	Berosus	6	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Dubiraphia	Dubiraphia vittata	4	6	SC/CG	"CN/50%, BU/50%"
Coleoptera	Haliphus	Haliphus	22	8	PH	:N,CM (la), SW,CM (ad
Coleoptera	Peltodytes	Peltodytes	2	8	PH	:N,CM (la), SW,CM (ad
Diptera	Ceratopogoninae	Culicoides	2	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	8	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Cladotanytarsus	8	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	20	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	8	7	CG/CF/PR	BU/CN/SP
Diptera	Dixella	Dixella	2		CG	SW
Diptera	Orthoclaadiinae	Corynoneura	8		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Psectrocladius	10		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Thienemanniella	4		CG/SC	SP/BU
Diptera	Tanypodinae	Ablabesmyia	4		PR	SP/BU
Diptera	Tanypodinae	Procladius	4		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	14	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	2	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	2	8	CG	BU
Hemiptera	Corixidae	Corixidae	4	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	8		unk	SW
Odonata	Coenagrionidae	Enallagma	110	7	PR	CM
Odonata	Libellulidae	Libellula	2	9	PR	SP
Odonata	Libellulidae	Sympetrum	2	9	PR	SP
Trichoptera	Limnephilus	Limnephilus	2	3	SH	CM/SP
Trombidiform	Acarina	Limnochares	2	5	PR	"SW/10%, CN/90%"

Montana Bioassessment Report

Waterbody Name: Otter Creek Site 16 for the Coal Tracts Study **Benthic Sample ID:** 17955
Station ID: OTTER_16f **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTR16fM-MAC-R
Site Classification: **Collection Date:** 10/15/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 603

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Ishnura	3			
Amphipoda	Hyaella	Hyaella	55	8	CG	SW/SP
Basommatop	Lymnaeidae	Pseudosuccinea columella	1	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	85	8	CG	CN
Basommatop	Planorbidae	Gyraulus deflectus	2	6	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	152	6	CG	CN
Coleoptera	Berosus	Berosus	1	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Dubiraphia	Dubiraphia vittata	2	6	SC/CG	"CN/50%, BU/50%"
Coleoptera	Haliphus	Haliphus	1	8	PH	"N,CM (la), SW,CM (ad)"
Diptera	Ceratopogoninae	Culicoides	4	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	33	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Cladotanytarsus	1	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	7	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	18	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Polypedilum	3	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Pseudochironomus	3	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	32	7	CG/CF/PR	BU/CN/SP
Diptera	Dasyheleinae	Dasyhelea	2		CG	BU/SW
Diptera	Orthoclaadiinae	Corynoneura	1		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Psectrocladius	2		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Thienemanniella	1		CG/SC	SP/BU
Diptera	Simuliidae	Simulium	10	6	CF	CN
Diptera	Tanypodinae	Ablabesmyia	5		PR	SP/BU
Diptera	Tanypodinae	Procladius	17		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	115	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	16	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	1	8	CG	BU
Hemiptera	Corixidae	Corixidae	1	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	2		unk	SW
Odonata	Coenagrionidae	Enallagma	15	7	PR	CM
Trichoptera	Cheumatopsyche	Cheumatopsyche	6	5	CF	CN
Trichoptera	Limnephilus	Limnephilus	1	3	SH	CM/SP
Trombidiform	Acarina	Hydrodroma	1	5	PR	"SW/10%, CN/90%"
Trombidiform	Acarina	Limnochares	2	5	PR	"SW/10%, CN/90%"
Trombidiform	Acarina	Pionidae	2	5	PR	"SW/10%, CN/90%"

Montana Bioassessment Report

Waterbody Name: Otter Creek Site 16 for the Coal Tracts Study **Benthic Sample ID:** 17956
Station ID: OTTER_16s **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTR16sM-MAC-R
Site Classification: **Collection Date:** 07/16/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 686

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Belostoma fluminea	4			
		Zavrelymia	2			
Amphipoda	Hyalella	Hyalella	162	8	CG	SW/SP
Basommatop	Lymnaeidae	Pseudosuccinea columella	4	6	CG	CN
Basommatop	Lymnaeidae	Stagnicola caperata	2	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	86	8	CG	CN
Basommatop	Physa_Physella	Physella gyrina	4	8	CG	CN
Basommatop	Planorbidae	Gyraulus deflectus	4	6	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	16	6	CG	CN
Coleoptera	Berosus	Berosus	16	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Dubiraphia	Dubiraphia vittata	2	6	SC/CG	"CN/50%, BU/50%"
Coleoptera	Haliphus	Haliphus	28	8	PH	:N,CM (la), SW,CM (ad
Coleoptera	Laccophilus	Laccophilus maculosus	2	5	PR	"CM (la), DI, SW (ad)"
Coleoptera	Peltodytes	Peltodytes	20	8	PH	:N,CM (la), SW,CM (ad
Diptera	Chironominae	Chironomus	6	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Cladotanytarsus	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	24	7	CG/CF/PR	BU/CN/SP
Diptera	Orthoclaadiinae	Corynoneura	1		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Psectrocladius	2		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Thienemanniella	2		CG/SC	SP/BU
Diptera	Tanypodinae	Ablabesmyia	2		PR	SP/BU
Diptera	Tanypodinae	Procladius	10		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	152	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	22	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	2	8	CG	BU
Hemiptera	Corixidae	Corixidae	2	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	8		unk	SW
Odonata	Aeshna	Aeshna palmata	6		PR	CM
Odonata	Coenagrionidae	Enallagma	72	7	PR	CM
Odonata	Lestes	Lestes	4	9	PR	SW
Odonata	Libellulidae	Libellula	4	9	PR	SP
Odonata	Libellulidae	Sympetrum	4	9	PR	SP
Trombidiform	Acarina	Hydrodroma	3	5	PR	"SW/10%, CN/90%"
Trombidiform	Acarina	Limnochares	4	5	PR	"SW/10%, CN/90%"

Montana Bioassessment Report

Waterbody Name: Home Creek Site 1A for the Coal Tracts Study **Benthic Sample ID:** 17957

Station ID: OTTER_1A

Rep. Num 0

Reference Status:

STORET Activity ID: OTTR1AM-MAC-R

Site Classification:

Collection Date: 05/16/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 639

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
Amphipoda	Hyaella	Hyaella	30	8	CG	SW/SP
Basommatop	Lymnaeidae	Stagnicola caperata	26	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	4	8	CG	CN
Basommatop	Physa_Physella	Physella gyrina	6	8	CG	CN
Coleoptera	Agabus	Agabus	26	5	PR	"CM(la), DI,SW(ad)"
Coleoptera	Berosus	Berosus	2	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Haliphus	Haliphus	6	8	PH	:N,CM (la), SW,CM (ad
Coleoptera	Hydrobius	Hydrobius	2			
Coleoptera	Hygrotus	Hygrotus	2		PR	"CM (la), DI, SW (ad)"
COLEOPTA	ILYBIUS	ILYBIUS	8	5	PR	
Diptera	Ceratopogoninae	Bezzia	4	6	PR/CG	SP/BU/SW
Diptera	Ceratopogoninae	Culicoides	14	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	8	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	24	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Endochironomus	4	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	14	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Stempellina	4	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	279	7	CG/CF/PR	BU/CN/SP
Diptera	Culicidae	Culex	8	10	CF/CG	SW
Diptera	Diamesinae	Potthastia	1	4	CG	sp
Diptera	Dixella	Dixella	4		CG	SW
Diptera	Ephydriidae	Ephydra	4	6	CG	BU
Diptera	Hedriodiscus/Odon	Odontomyia	1		CG	SP
Diptera	Orthoclaadiinae	Corynoneura	2		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Psectrocladius	40		CG/SC	SP/BU
Diptera	Tabanidae	Tabanus	4	10	PR	SP
Diptera	Tanypodinae	Procladius	28		PR	SP/BU
Diptera	Tanypodinae	Thienemannimyia Gr.	4		PR	SP/BU
Diptera	Tipula	Tipula	2	4	SH	BU
Ephemeropte	Caenis	Caenis latipennis	4	8	CG	"SP/75%, CM/90%"
Non-Insect ta	Nematoda	Nematoda	4	5	unk	BU
Non-Insect ta	Ostracoda	Ostracoda	44		unk	SW
Odonata	Aeshna	Aeshna	8		PR	CM
Odonata	Lestes	Lestes	6	9	PR	SW
Trichoptera	Limnephilus	Limnephilus	12	3	SH	CM/SP

Montana Bioassessment Report

Waterbody Name: Home Creek Site 1A for the Coal Tracts Study **Benthic Sample ID:** 17958
Station ID: OTTER_1Af **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTR1AfM-MAC-R
Site Classification: **Collection Date:** 10/14/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 617

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Proezzia	11			
Amphipoda	Hyaella	Hyaella	27	8	CG	SW/SP
Basommatop	Lymnaeidae	Stagnicola caperata	52	6	CG	CN
Basommatop	Lymnaeidae	Stagnicola elodes	3	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	32	8	CG	CN
Basommatop	Physa_Physella	Physella gyrina	8	8	CG	CN
Coleoptera	Agabus	Agabus	1	5	PR	"CM(la), DI,SW(ad)"
Coleoptera	Berosus	Berosus	1	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Colymbetes	Colymbetes	2		PR	"CM(la), DI,SW(ad)"
Coleoptera	Haliphus	Haliphus	1	8	PH	:N,CM (la), SW,CM (ad
Coleoptera	Hydrobius	Hydrobius	1			
Diptera	Ceratopogoninae	Culicoides	2	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	135	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	1	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	25	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Polypedilum	1	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Stempellina	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	89	7	CG/CF/PR	BU/CN/SP
Diptera	Dasyheleinae	Dasyhelea	43		CG	BU/SW
Diptera	Ephydriidae	Ephydra	1	6	CG	BU
Diptera	Hedriodiscus/Odon	Odontomyia	1		CG	SP
Diptera	Orthoclaadiinae	Psectrocladius	121		CG/SC	SP/BU
Diptera	Pericoma/Telmatos	Pericoma	2	4	CG	BU
Diptera	Simuliidae	Simulium	4	6	CF	CN
Diptera	Tabanidae	Tabanus	1	10	PR	SP
Diptera	Tanypodinae	Ablabesmyia	1		PR	SP/BU
Diptera	Tanypodinae	Procladius	6		PR	SP/BU
Haplotaxida	Oligochaeta	Tubificidae	10	8	CG	BU
Hemiptera	Notonectidae	Notonecta	7	7	PR	SW
Non-Insect ta	Ostracoda	Ostracoda	2		unk	SW
Odonata	Aeshna	Aeshna palmata	1		PR	CM
Odonata	Amphiagrion	Amphiagrion abbreviatum	2	7	PR	CM
Odonata	Coenagrionidae	Enallagma	18	7	PR	CM
Odonata	Lestes	Lestes	1	9	PR	SW
Trichoptera	Limnephilus	Limnephilus	2	3	SH	CM/SP

Montana Bioassessment Report

Waterbody Name: Home Creek Site 1A for the Coal Tracts Study **Benthic Sample ID:** 17959

Station ID: OTTER_1As

Rep. Num 0

Reference Status:

STORET Activity ID: OTR1AsM-MAC-R

Site Classification:

Collection Date: 07/17/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 626

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Liodessus affinis	1			
Amphipoda	Hyaella	Hyaella	22	8	CG	SW/SP
Basommatop	Lymnaeidae	Stagnicola caperata	11	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	32	8	CG	CN
COLEOPTE	ANACAENA	ANACAENA	1	5	PR/CG	
Coleoptera	Berosus	Berosus	5	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Haliphus	Haliphus	22	8	PH	:N,CM (la), SW,CM (ad
Coleoptera	Helophorus	Helophorus	1	5	SH	CM
Coleoptera	Hydrobius	Hydrobius	12			
Coleoptera	Hydroporus	Hydroporus	2		PR	"CM (la), DI, SW (ad)"
Coleoptera	Tropisternus	Tropisternus	1		PR	N,SP,CM(la), DI,SW(ac
Diptera	Ceratopogoninae	Bezzia	1	6	PR/CG	SP/BU/SW
Diptera	Ceratopogoninae	Culicoides	1	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	198	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	1	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	4	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	165	7	CG/CF/PR	BU/CN/SP
Diptera	Culicidae	Culex	3	10	CF/CG	SW
Diptera	Dasyheleinae	Dasyhelea	7		CG	BU/SW
Diptera	Ephydriidae	Ephydra	5	6	CG	BU
Diptera	Orthocladinae	Psectrocladius	65		CG/SC	SP/BU
Diptera	Tabanidae	Tabanus	1	10	PR	SP
Diptera	Tanypodinae	Procladius	1		PR	SP/BU
Haplotaxida	Oligochaeta	Tubificidae	4	8	CG	BU
Hemiptera	Corixidae	Corixidae	15	9	PH/PR	SW
Hemiptera	Naucoridae	Ambrysus mormon	1	3	PR	CM
Non-Insect ta	Ostracoda	Ostracoda	5		unk	SW
Odonata	Aeshna	Aeshna palmata	27		PR	CM
Odonata	Coenagrionidae	Enallagma	5	7	PR	CM
Odonata	Lestes	Lestes	5	9	PR	SW
Odonata	Libellulidae	Sympetrum	1	9	PR	SP
Trichoptera	Limnephilus	Limnephilus	1	3	SH	CM/SP

Montana Bioassessment Report

Waterbody Name: Otter Creek Site 22 for the Coal Tracts Study **Benthic Sample ID:** 17960
Station ID: OTTER_22 **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTTR22M-MAC-R
Site Classification: **Collection Date:** 05/17/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 628

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Helisoma trivolvis	5			
Amphipoda	Hyaella	Hyaella	5	8	CG	SW/SP
Basommatop	Lymnaeidae	Pseudosuccinea columella	2	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	270	8	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	51	6	CG	CN
BASOMMAT	Planorbidae	HELISOMA ANCEPS	21	6	CG	CN
Coleoptera	Halipus	Halipus	3	8	PH	:N,CM (la), SW,CM (ad
Diptera	Chironominae	Chironomus	48	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	7	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	122	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Microtendipes pedellus Gr.	8	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	7	7	CG/CF/PR	BU/CN/SP
Diptera	Orthoclaadiinae	Psectrocladius	3		CG/SC	SP/BU
Diptera	Tanypodinae	Ablabesmyia	2		PR	SP/BU
Diptera	Tanypodinae	Procladius	22		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	5	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	6	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	2	8	CG	BU
Hemiptera	Corixidae	Corixidae	3	9	PH/PR	SW
Odonata	Aeshna	Aeshna palmata	1		PR	CM
Odonata	Coenagrionidae	Enallagma	27	7	PR	CM
Rhynchobdell	Glossiphoniidae	Glossiphonia complanata	1	9	PR	SW
Rhynchobdell	Glossiphoniidae	Helobdella stagnalis	1	9	PR	SW
Trichoptera	Limnephilus	Limnephilus	6	3	SH	CM/SP

Montana Bioassessment Report

Waterbody Name: Otter Creek Site 22 for the Coal Tracts Study **Benthic Sample ID:** 17961
Station ID: OTTER_22f **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTR22fM-MAC-R
Site Classification: **Collection Date:** 10/14/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 725

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Helisoma trivolvis	1			
		Ishnura	3			
Amphipoda	Hyalella	Hyalella	155	8	CG	SW/SP
Basommatop	Physa_Physella	Physella acuta	120	8	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	20	6	CG	CN
BASOMMAT	Planorbidae	HELISOMA ANCEPS	4	6	CG	CN
Coleoptera	Berosus	Berosus	11	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Coptotomus	Coptotomus longulus	1	5	PR	"CM (la), DI, SW (ad)"
Coleoptera	Haliphus	Haliphus	15	8	PH	"N,CM (la), SW,CM (ad)"
Coleoptera	Hydroporus	Hydroporus	1		PR	"CM (la), DI, SW (ad)"
Coleoptera	Rhantus	Rhantus	2		PR	"CM (la), DI, SW (ad)"
Diptera	Ceratopogoninae	Culicoides	2	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	1	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	15	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	19	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Pseudochironomus	7	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	29	7	CG/CF/PR	BU/CN/SP
Diptera	Orthoclaadiinae	Cricotopus	1		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Psectrocladius	1		CG/SC	SP/BU
Diptera	Tanypodinae	Ablabesmyia	3		PR	SP/BU
Diptera	Tanypodinae	Procladius	45		PR	SP/BU
Diptera	Tanypodinae	Thienemannimyia Gr.	5		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	145	8	CG	"SP/75%, CM/90%"
Ephemeropte	Caenis	Caenis youngi	6	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	7	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	3	8	CG	BU
Hemiptera	Corixidae	Corixidae	10	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	11		unk	SW
Odonata	Aeshna	Aeshna palmata	1		PR	CM
Odonata	Coenagrionidae	Enallagma	70	7	PR	CM
Odonata	Lestes	Lestes	1	9	PR	SW
Odonata	Libellulidae	Libellula	4	9	PR	SP
Trichoptera	Cheumatopsyche	Cheumatopsyche	2	5	CF	CN
Trichoptera	Limnephilus	Limnephilus	2	3	SH	CM/SP
Trombidiform	Acarina	Limnochares	2	5	PR	"SW/10%, CN/90%"

Montana Bioassessment Report

Waterbody Name: Otter Creek Site 22 for the Coal Tracts Study **Benthic Sample ID:** 17962
Station ID: OTTER_22s **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTR22sM-MAC-R
Site Classification: **Collection Date:** 07/14/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 879

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Helisoma trivolvis	1			
Amphipoda	Hyaella	Hyaella	95	8	CG	SW/SP
Basommatop	Lymnaeidae	Pseudosuccinea columella	1	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	225	8	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	18	6	CG	CN
BASOMMAT	Planorbidae	HELISOMA ANCEPS	3	6	CG	CN
Coleoptera	Berosus	Berosus	2	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Colymbetes	Colymbetes	1		PR	"CM(la), DI,SW(ad)"
Coleoptera	Haliphus	Haliphus	15	8	PH	:N,CM (la), SW,CM (ad
Coleoptera	Hydroporus	Hydroporus	8		PR	"CM (la), DI, SW (ad)"
Coleoptera	Peltodytes	Peltodytes	3	8	PH	:N,CM (la), SW,CM (ad
Diptera	Chironominae	Chironomus	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	5	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	15	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Microtendipes pedellus Gr.	1	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	28	7	CG/CF/PR	BU/CN/SP
Diptera	Dixella	Dixella	1		CG	SW
Diptera	Orthoclaadiinae	Cricotopus	7		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Psectrocladius	7		CG/SC	SP/BU
Diptera	Simuliidae	Simulium	1	6	CF	CN
Diptera	Tanypodinae	Ablabesmyia	23		PR	SP/BU
Diptera	Tanypodinae	Procladius	17		PR	SP/BU
Diptera	Tanypodinae	Thienemannimyia Gr.	8		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	288	8	CG	"SP/75%, CM/90%"
Ephemeropte	Caenis	Caenis youngi	3	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	25	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	1	8	CG	BU
Hemiptera	Corixidae	Corixidae	5	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	11		unk	SW
Odonata	Aeshna	Aeshna palmata	3		PR	CM
Odonata	Coenagrionidae	Enallagma	45	7	PR	CM
Odonata	Lestes	Lestes	5	9	PR	SW
Rhynchoabdell	Glossiphoniidae	Glossiphonia complanata	2	9	PR	SW
Rhynchoabdell	Glossiphoniidae	Helobdella stagnalis	2	9	PR	SW
Trichoptera	Limnephilus	Limnephilus	2	3	SH	CM/SP

Montana Bioassessment Report

Waterbody Name: Tenmile Creek Site 23 for the Coal Tracts Study **Benthic Sample ID:** 17963

Station ID: Otter_23

Rep. Num 0

Reference Status:

STORET Activity ID: OTTR23M-MAC-R

Site Classification:

Collection Date: 05/17/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 605

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Agabus sp. 2	6			
Coleoptera	Agabus	Agabus	33	5	PR	"CM(la), DI,SW(ad)"
Coleoptera	Colymbetes	Colymbetes	4		PR	"CM(la), DI,SW(ad)"
Coleoptera	Coptotomus	Coptotomus longulus	2	5	PR	"CM (la), DI, SW (ad)"
Coleoptera	Rhantus	Rhantus	2		PR	"CM (la), DI, SW (ad)"
Diptera	Chironominae	Chironomus	26	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	265	7	CG/CF/PR	BU/CN/SP
Diptera	Culicidae	Aedes	44	10	CF/CG	SW
Diptera	Culicidae	Culex	98	10	CF/CG	SW
Diptera	Ephydriidae	Ephydriidae	5	6	CG	BU
Diptera	Orthoclaadiinae	Psectrocladius	64		CG/SC	SP/BU
Diptera	Sciomyzidae	Hedria	8		PR	BU
Hemiptera	Corixidae	Sigara	4	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	42		unk	SW
Trichoptera	Limnephilus	Limnephilus	2	3	SH	CM/SP

Montana Bioassessment Report

Waterbody Name: Tenmile Creek Site 23 for the Coal Tracts Study **Benthic Sample ID:** 17964
Station ID: Otter_23f **Rep. Num** 0
Reference Status: **STORET Activity ID:** OTR23fM-MAC-R
Site Classification: **Collection Date:** 10/15/2011
Latitude: **Collection Method:** MAC-R-500
Longitude: **Total Number of Individuals in Sample:** 617

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
Coleoptera	Agabus	Agabus	33	5	PR	"CM(la), DI,SW(ad)"
Coleoptera	Colymbetes	Colymbetes	1		PR	"CM(la), DI,SW(ad)"
Coleoptera	Coptotomus	Coptotomus longulus	1	5	PR	"CM (la), DI, SW (ad)"
Coleoptera	Rhantus	Rhantus	4		PR	"CM (la), DI, SW (ad)"
Diptera	Chironominae	Chironomus	106	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	177	7	CG/CF/PR	BU/CN/SP
Diptera	Culicidae	Culex	23	10	CF/CG	SW
Diptera	Ephydriidae	Ephydra	5	6	CG	BU
Diptera	Ephydriidae	Ephydriidae	13	6	CG	BU
Diptera	Orthoclaadiinae	Psectrocladius	122		CG/SC	SP/BU
Hemiptera	Corixidae	Corixidae	22	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	98		unk	SW
Odonata	Coenagrionidae	Enallagma	12	7	PR	CM

Montana Bioassessment Report

Waterbody Name: Tenmile Creek Site 23 for the Coal Tracts Study **Benthic Sample ID:** 17965

Station ID: Otter_23s

Rep. Num 0

Reference Status:

STORET Activity ID: OTR23sM-MAC-R

Site Classification:

Collection Date: 07/17/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 515

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Agabus sp. 2	12			
Coleoptera	Agabus	Agabus	50	5	PR	"CM(la), DI,SW(ad)"
Coleoptera	Colymbetes	Colymbetes	4		PR	"CM(la), DI,SW(ad)"
Coleoptera	Coptotomus	Coptotomus longulus	2	5	PR	"CM (la), DI, SW (ad)"
Coleoptera	Rhantus	Rhantus	2		PR	"CM (la), DI, SW (ad)"
Diptera	Chironominae	Chironomus	56	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	137	7	CG/CF/PR	BU/CN/SP
Diptera	Culicidae	Culex	67	10	CF/CG	SW
Diptera	Ephydriidae	Ephydriidae	13	6	CG	BU
Diptera	Orthoclaadiinae	Psectrocladius	90		CG/SC	SP/BU
Hemiptera	Corixidae	Sigara	4	9	PH/PR	SW
Non-Insect ta	Ostracoda	Ostracoda	78		unk	SW

Montana Bioassessment Report

Waterbody Name: Otter Creek Site ASJT for the Coal Tracts Study **Benthic Sample ID:** 17966

Station ID: OTTER_JT

Rep. Num 0

Reference Status:

STORET Activity ID: OTTRJTM-MAC-R

Site Classification:

Collection Date: 05/17/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 610

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
Amphipoda	Hyaella	Hyaella	70	8	CG	SW/SP
Basommatop	Lymnaeidae	Pseudosuccinea columella	2	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	62	8	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	2	6	CG	CN
Coleoptera	Dubiraphia	Dubiraphia vittata	58	6	SC/CG	"CN/50%, BU/50%"
Diptera	Ceratopogoninae	Bezzia	2	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	48	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Cladotanytarsus	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Cryptochironomus	10	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	28	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Endochironomus	10	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	5	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Parachironomus	5	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Phaenopsectra	12	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Polypedilum	8	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	8	7	CG/CF/PR	BU/CN/SP
Diptera	Ephydriidae	Ephydra	2	6	CG	BU
Diptera	Orthoclaadiinae	Cricotopus	2		CG/SC	SP/BU
Diptera	Simuliidae	Simulium	10	6	CF	CN
Diptera	Tanypodinae	Ablabesmyia	32		PR	SP/BU
Diptera	Tanypodinae	Procladius	30		PR	SP/BU
Diptera	Tanypodinae	Thienemannimyia Gr.	2		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	75	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	12	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	22	8	CG	BU
Hemiptera	Corixidae	Corixidae	38	9	PH/PR	SW
Odonata	Coenagrionidae	Enallagma	8	7	PR	CM
Trichoptera	Cheumatopsyche	Cheumatopsyche	2	5	CF	CN
Trichoptera	Hydropsyche_Cera	Hydropsyche	2	5	CF	CN
Trichoptera	Hydropsyche_Cera	Hydropsyche morosa gr.	2	5	CF	CN
Trichoptera	Hydroptila	Hydroptila	5	6	PH	CN
Trichoptera	Limnephilus	Limnephilus	8	3	SH	CM/SP
Veneroida	Pisidiidae	Pisidium compressum	8	8	CF	BU
Veneroida	Pisidiidae	Sphaerium simile	18	8	CF	BU

Montana Bioassessment Report

Waterbody Name: Otter Creek Site ASJT for the Coal Tracts Study **Benthic Sample ID:** 17967

Station ID: OTTER_JTf

Rep. Num 0

Reference Status:

STORET Activity ID: OTRJTfM-MAC-R

Site Classification:

Collection Date: 10/15/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 606

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Helisoma trivolvis	1			
		Sialis veleta	1			
Amphipoda	Hyalella	Hyalella	5	8	CG	SW/SP
Basommatop	Lymnaeidae	Pseudosuccinea columella	4	6	CG	CN
Basommatop	Physa_Physella	Physella acuta	45	8	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	8	6	CG	CN
BASOMMAT	Planorbidae	HELISOMA ANCEPS	2	6	CG	CN
Coleoptera	Berosus	Berosus	3	5	PH	"CM(la), DI,SW(ad)"
Coleoptera	Dubiraphia	Dubiraphia vittata	190	6	SC/CG	"CN/50%, BU/50%"
Coleoptera	Haliphus	Haliphus	1	8	PH	:N,CM (la), SW,CM (ad
Coleoptera	Stenelmis	Stenelmis	1	5	SC/CG	"CN/50%, BU/50%"
Diptera	Ceratopogoninae	Culicoides	20	6	PR/CG	SP/BU/SW
Diptera	Ceratopogoninae	Probezzia	8	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	27	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Cryptochironomus	5	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	15	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Endochironomus	1	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Glyptotendipes	39	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Polypedilum	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	41	7	CG/CF/PR	BU/CN/SP
Diptera	Dasyheleinae	Dasyhelea	15		CG	BU/SW
Diptera	Ephydriidae	Ephydra	1	6	CG	BU
Diptera	Orthoclaadiinae	Cricotopus	11		CG/SC	SP/BU
Diptera	Orthoclaadiinae	Psectrocladius	60		CG/SC	SP/BU
Diptera	Pericoma/Telmatos	Pericoma	12	4	CG	BU
Diptera	Simuliidae	Simulium	2	6	CF	CN
Diptera	Tanypodinae	Ablabesmyia	1		PR	SP/BU
Diptera	Tanypodinae	Procladius	21		PR	SP/BU
Diptera	Tanypodinae	Thienemannimyia Gr.	1		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	40	8	CG	"SP/75%, CM/90%"
Ephemeropte	Callibaetis	Callibaetis	1	9	CG	"SW/10%, CN/90%"
Haplotaxida	Oligochaeta	Tubificidae	1	8	CG	BU
Hemiptera	Corixidae	Corixidae	7	9	PH/PR	SW
Odonata	Coenagrionidae	Enallagma	3	7	PR	CM
Odonata	Gomphidae	Ophiogomphus severus	2	2	PR	BU

Montana Bioassessment Report

Waterbody Name: Otter Creek Site ASJT for the Coal Tracts Study **Benthic Sample ID:** 17967

Station ID: OTTER_JTf

Rep. Num 0

Reference Status:

STORET Activity ID: OTRJTfM-MAC-R

Site Classification:

Collection Date: 10/15/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 606

Odonata	Libellulidae	Libellula	3	9	PR	SP
Trichoptera	Cheumatopsyche	Cheumatopsyche	1	5	CF	CN
Trichoptera	Hydroptila	Hydroptila	1	6	PH	CN
Trichoptera	Limnephilus	Limnephilus	2	3	SH	CM/SP
Trichoptera	Phryganeidae	Ptilostomis	1	4	SH	CM
Veneroida	Pisidiidae	Sphaerium simile	1	8	CF	BU

Montana Bioassessment Report

Waterbody Name: Otter Creek Site ASJT for the Coal Tracts Study **Benthic Sample ID:** 17968

Station ID: OTTER_JTs

Rep. Num 0

Reference Status:

STORET Activity ID: OTRJTsM-MAC-R

Site Classification:

Collection Date: 07/18/2011

Latitude:

Collection Method: MAC-R-500

Longitude:

Total Number of Individuals in Sample: 605

Sample Taxa List

<i>Order:</i>	<i>OTU name:</i>	<i>FinalID:</i>	<i>Individuals</i>	<i>Tol Val:</i>	<i>FFG:</i>	<i>Habit:</i>
		Helisoma trivolvis	4			
Amphipoda	Hyaella	Hyaella	172	8	CG	SW/SP
Basommatop	Physa_Physella	Physella acuta	50	8	CG	CN
Basommatop	Planorbidae	Gyraulus parvus	2	6	CG	CN
Coleoptera	Colymbetes	Colymbetes	4		PR	"CM(la), DI,SW(ad)"
Coleoptera	Dubiraphia	Dubiraphia vittata	12	6	SC/CG	"CN/50%, BU/50%"
Coleoptera	Stenelmis	Stenelmis	2	5	SC/CG	"CN/50%, BU/50%"
Diptera	Ceratopogoninae	Bezzia	2	6	PR/CG	SP/BU/SW
Diptera	Chironominae	Chironomus	8	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Cladotanytarsus	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Dicrotendipes	6	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Phaenopsectra	2	7	CG/CF/PR	BU/CN/SP
Diptera	Chironominae	Tanytarsus	36	7	CG/CF/PR	BU/CN/SP
Diptera	Orthoclaadiinae	Cricotopus	2		CG/SC	SP/BU
Diptera	Simuliidae	Simulium	45	6	CF	CN
Diptera	Tanypodinae	Ablabesmyia	28		PR	SP/BU
Diptera	Tanypodinae	Procladius	98		PR	SP/BU
Diptera	Tanypodinae	Thienemannimyia Gr.	2		PR	SP/BU
Ephemeropte	Caenis	Caenis latipennis	98	8	CG	"SP/75%, CM/90%"
Haplotaxida	Oligochaeta	Tubificidae	2	8	CG	BU
Hemiptera	Corixidae	Corixidae	12	9	PH/PR	SW
Hemiptera	Naucoridae	Ambrysus mormon	2	3	PR	CM
Heterostroph	Valvata	Valvata lewisi	2	3	SC	CM
Trichoptera	Cheumatopsyche	Cheumatopsyche	2	5	CF	CN
Trichoptera	Hydropsyche_Cera	Hydropsyche	4	5	CF	CN
Trichoptera	Hydropsyche_Cera	Hydropsyche morosa gr.	2	5	CF	CN
Trichoptera	Hydroptila	Hydroptila	2	6	PH	CN
Trichoptera	Limnephilus	Limnephilus	2	3	SH	CM/SP

Appendix C . Habitat and Water Quality Parameters measured for the Otter Creek sites visited. NA = not visited or sampled during this visit, dry.

	OTTER_23			OTTER_22			OTTER_16			OTTER_3m			OTTER_2			OTTER_JT			OTTER_1A		
	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct	May	Jul	Oct
Water Temp ©	17.4	30.8	3.8	12.3	28.5	7.5	12.9	24.6	8.8	14.5	na	na	na	26.6	9.2	13.1	26.8	10.7	18.6	27.8	9.1
Conductivity (µs/cm)	1880	>4,000	>4,000	3660	3790	3590	>4,000	3920	3790	170	na	na	na	3890	3700	3860	3608	3540	1950	>4,000	3209
pH	8.2	8.4	8.4	8.2	8.2	8.1	7.6	7.8	8	7.8	na	na	na	8.2	8.1	8.1	8.1	8.1	8.2	8.2	8.2
PFC	FAR	FAR	FAR	PFC	PFC	PFC	PFC	PFC	FAR	FAR	FAR	FAR	na	FAR	FAR	PFC	PFC	PFC	FAR	FAR	FAR
BLM HBI	16	17	17	22	22	22	19	19	18	17	18	18	na	17	17	19	19	19	11	12	13
Avg wetted width (m)	6.2	0.6	0.6	8.7	8.2	8.6	11.9	12.1	11.2	1.8	na	na	na	4.2	2.8	5.0	4.8	4.6	2.4	2.1	2.0
Avg Left CHD (cm)	25	2	2	71	67	71	103	104	93	2	na	na	na	30	22	53	43	36	22	28	31
Avg Center CHD (cm)	32	3	3	76	97	76	136	128	106	2	na	na	na	42	30	54	44	40	30	33	42
Avg Right CHD (cm)	27	2	2	70	66	70	95	92	86	2	na	na	na	25	20	55	41	36	18	25	29
% Fines in Reach	98	100	100	92	84	92	97	97	89	100	100	100	na	50	60	70	67	63	100	100	98
% Gravel Reach	2	0	0	4	8	4	3	3	8	0	0	0	na	35	30	20	26	34	1	1	3
% Cobble Reach	0	0	0	5	9	5	1	1	5	0	0	0	na	15	10	11	8	3	0	0	0
Livestock Use (CPI)	30	11	9	55	36	13	18	7	46	35	44	24	na	12	5	18	22	19	55	36	28
Avg. Riparian Shade	0	5	0	0	10	0	5	10	10	10	40	20	na	10	10	10	20	10	10	30	20