

\* \* D R A F T \* \*  
August 27, 1990

**GRASSLAND AND SHRUBLAND COMMUNITIES ON USDI  
BUREAU OF LAND MANAGEMENT LANDS IN MONTANA**

A Proposal by

Robert L. DeVelice  
Montana Natural Heritage Program  
1515 East 6th Avenue, Helena, MT 59620  
406-444-3009

**MISSION STATEMENT**

*The proposed Montana Natural Heritage Program's (MTNHP) grasslands/shrublands ecological classification project (GSCP) on Bureau of Land Management (BLM) lands is a working component of The Nature Conservancy's ecology program in the western United States. The Nature Conservancy program provides key information on plant communities to be used for conservation planning, management, research, and monitoring. Although grasslands and shrublands cover over 75% of the Montana landscape, an exhaustive review of existing information (MTNHP 1990) showed that they are the least documented vegetation types of the state. Therefore, the GSCP is designed to complete the classification over the whole ecological space and to conduct regional correlations of existing classifications. The information provided by the project will be the basis for programs to model the effects of management, global changes, and other variables on the vegetation types and diversity patterns, and their implications for further management and conservation planning. The GSCP reflects general conservation planning, management, and research issues. The project will continue to focus on strong collaborative work with the various state and federal agencies (BLM, USFS, BIA, DOD) and other institutions (e.g. Montana universities) in order to contribute to the development of a tightly integrated state-wide classification system.*

U9PDEV62MTUS

## *SUMMARY*

The proposed research would develop a classification of plant communities (primarily grasslands and shrublands) on BLM lands throughout Montana. Such a classification will be useful in identifying sensitive communities and natural areas where environmental impact from resource manipulations (e.g., domestic grazing; mining) would be minimized. Additionally, the classification would provide a reference system for baseline monitoring of environmental impacts and vegetation recovery and would provide an ecological basis for categorizing environmental variation. The classification would be based on the integration and synthesis of both existing data and newly sampled plots.

## *INTRODUCTION*

A more complete assessment of sensitive plant species and communities would complement the BLM's existing range site characterization process. The proposed work would provide a classification of Montana grassland and shrubland communities on BLM lands that could be used to assess ecosystem sensitivities and potential responses to perturbation.

In addition to the environmental impact assessment applications, the proposed work represents a step towards developing a comprehensive classification of Montana plant communities that will provide land managers and scientists a state-wide perspective of community variation. Such a perspective is invaluable towards making


sound management prescriptions and predictions, designing and interpreting experiments, and identifying areas of critical importance for conservation.

### *PREVIOUS RESEARCH*

Grasslands and shrublands cover over 75 percent of the Montana landscape yet are the most poorly described vegetation types of the state. In contrast, the classification of the forest types of Montana is largely complete. Figure 1 highlights both the vast expanse of Montana grasslands and the sparseness of available detailed community characterizations, particularly in the eastern half of the state. Filling this gap constitutes the primary goal of the proposed project. It is a major and necessary step towards developing a comprehensive classification of all Montana plant communities.

To date, studies characterizing grassland and shrubland communities of Montana have been of limited geographical and ecological scope. The most extensive existing studies include Mueggler and Stewart's (1980) in western Montana, Jorgensen's (1979) and Harvey's (1982) studies in east-central Montana, and Hansen and Hoffman's (1985) work in southeastern Montana. A dissertation describing grassland communities of south-central Montana is currently being completed (by Steve Harvey, Montana State University, Bozeman).

The MTNHP conducted fieldwork (largely on BLM lands) in northcentral and northeastern Montana during the summer of 1990. Vegetation and site data were collected from 124 sites representing a wide-range of grassland/shrubland environments



\* \* D R A F T \* \*  
August 27, 1990

and species assemblages. These data are currently being analyzed and a report describing the communities and their environmental relations will be completed by November 1990. At present, 22 grassland/shrubland community types have been tentatively identified based on the 1990 data (Table 1).

Relevant grassland/shrubland classifications from adjacent states and provinces include: Coupland (1950; 1961), Girard et al. (1989), Hansen et al. (1984), Whitman and Hanson (1939), and Jones (1989).

The classification of Montana's upland forest plant associations provided by Pfister et al. (1977) has been refined and completed by the work of Cooper and Pfister (1981; 1985), Hansen and Hoffman (1985), Hoffman and Hansen (1981), and Roberts (1980). Classifications of Montana riparian sites (including forested, shrub dominated, and herb dominated communities) are nearing completion (personal communication, Paul Hansen, Montana Riparian Association, School of Forestry, University of Montana, Missoula).

## ***SCOPE OF PROPOSED RESEARCH***

### ***Objectives***

The proposed research would provide the following:

- 1) A classification of grassland and shrubland communities on BLM lands in Montana. In addition, undescribed or poorly-known forest and woodland communities will be sampled and classified as good examples are found.

\* \* D R A F T \* \*

August 27, 1990

Table 1. Community types preliminarily identified in northeastern/northcentral Montana based on 1990 MTNHP fieldwork.

### Shrublands

*Artemisia cana*/*Agropyron smithii*

*Artemisia longifolia*

*Artemisia tridentata*/*Agropyron smithii*

*Artemisia tridentata*/*Agropyron spicatum*

*Artemisia tridentata*-*Atriplex confertifolia*

*Atriplex gardneri*

*Juniperun horizontalis*/*Andropogon scoparius*

*Rhus aromatica*/*Agropyron spicatum*

*Sarcobatus vermiculatus*/*Agropyron smithii*

*Sarcobatus vermiculatus*-*Artemisia tridentata*

### Grasslands

*Agropyron smithii*

*Agropyron smithii*-*Bouteloua gracilis*

*Agropyron smithii*-*Carex filifolia*

*Agropyron smithii*-*Stipa viridula*

*Agropyron spicatum*-*Agropyron smithii*

*Agropyron spicatum*-*Bouteloua gracilis*

*Agropyron spicatum*-*Carex filifolia*

*Agropyron spicatum*-*Poa sandbergii*

*Andropogon scoparius*-*Muhlenbergia cuspidata*

*Calamovilfa longifolia*-*Agropyron smithii*

*Stipa comata*-*Bouteloua gracilis*

*Stipa comata*-*Carex filifolia*

\* \* D R A F T \* \*  
August 27, 1990

- 2) Descriptions of the floristic, geographic, and physiographic features of each community.
- 3) Relevant information regarding such features as successional trends, site productivity potential, and potential sensitivities to a range of perturbations.
- 4) Descriptions of the location and characteristics of high priority communities and sites for conservation.

*Expected Significance*

The BLM would find a classification of grassland and shrubland communities useful in stratifying vegetation/environmental variation to assess management options and results. Additionally, the classification will assist in minimizing impacts from intensive management by identifying sensitive plant communities.

One function of the MTNHP is the development of a statewide database of plant community occurrences. A major limitation is the current lack of a comprehensive grassland/shrubland community classification. The proposed work would be a significant step towards achieving such a comprehensive classification.

\* \* D R A F T \* \*  
August 27, 1990

*Relation to Longer-Term Goals*

The BLM would find a state-wide grassland/shrubland community classification useful as a reference system for baseline monitoring of ecological impacts and assessing vegetation recovery following disturbance. Additionally, the classification would be useful as a tool for predicting long-term vegetation responses to management activities.

One of the functions of the MTNHP is to provide information to The Nature Conservancy regarding communities and sites for conservation. A statewide classification of Montana vegetation (including grasslands/shrublands) is necessary to define and identify key elements and sites for potential long-term preservation. Similarly, the BLM could use the classification for the identification and design of Areas of Critical Environmental Concern (ACEC's). For example, by using vegetation community as a data theme within a Geographic Information System (GIS), preserves could be efficiently designed to contain the range of community/environmental patterns of an area. Such capability is invaluable towards global biodiversity preservation.

*Relation to Present State of Knowledge*

Existing classifications and data inadequately describe the grassland and shrubland communities of Montana. Major additional field

\* \* D R A F T \* \*

August 27, 1990

sampling (as proposed here) is necessary before a comprehensive grassland/shrubland community classification can be developed.

## *METHODS*

Prior to initiating field sampling, data from previous research and range site data will be used to develop a preliminary classification of Montana's grassland/shrubland communities. A preliminary listing of grassland and shrubland communities for all of eastern Montana has already been compiled by the MTNHP (Table 2). This list will be refined and reevaluated in reference to all of Montana to provide an initial perspective of community species composition and diversity.

To maximize the efficiency in sampling the range of vegetation and environmental variation, field samples will be subjectively selected using the "gradsect" method described and evaluated by Gillison and Brewer (1985) and applied by Austin and Heyligers (1989). The method involves preferential sampling along local transects following the maximum perceived environmental gradients. Representation of the range of moisture, temperature, radiation, and soil nutrient regimes on BLM lands in Montana will be emphasized.

In the office, a regular grid with 0.01° (approximately 0.3 mi<sup>2</sup>) spacing in both longitudinal and latitudinal directions will be overlaid on 7.5 minute USGS topographic maps covering BLM lands. Land-use maps will be used to determine which grid cells fall on farmed and urban/industrial areas and these cells will not be examined further. The following will be recorded for each of the remaining grid cells:

Table 2. Relationship of preliminary eastern Montana grassland plant communities identified by the MTNHP to previous classifications. Riparian and tundra communities are not included.

Other Ard. '73 Cou. '80	Brown 1971	Mackie 1970	Dusak 1971	Jorgens. 1979	Harvey 1982	Cul/Skow 1982	Culwell 1985	Progers 1978	Hansen 1965	Mun./Ste. 1980	MTNHP
											Agsm
						Agsm/Bogr/ Buda		Agsm/Bogr			Agsm-Bogr
							Elem/Bogr/ Calf		Agsm/Calf		Agsm-Calf
					Agsm/Stvl		Stcv/Stvl	Stvl-Agsm/ Bogr			Agsm-Stvl
				Agsp/Agsm						Agsp/Agsm	Agsp-Agsm <sup>2</sup>
									Agsp/Boca		Agsp-Boca
					Agsp/Bogr			Agsp/Bogr		Agsp/Bogr	Agsp-Bogr <sup>2</sup>
									Agsp/Calf		Agsp-Calf <sup>2</sup>
								Mucu/Agsp			Agsp-Mucu <sup>2</sup>
										Agsp/Poss	Agsp-Poss <sup>2</sup>
+/- Tay. '76											Ange
Tay. '76											Ange-Anac
								Anac-Agsp			Anac-Agsp
							Calo/Anac	Anac-Calo	Anac/Calf		Anac-Calf
						Anac/Cals/ Bocu/Calo					Anac-Cals
				Mucu/Anac							Anac-Mucu
Bog. '84											Calo-Agsm
+/- Ros. '73											Calo-Anac
							Calo/Calf				Calo-Calf
									Calo/Cals		Calo-Cals

Table 2. (continued)

other	Brown 1971	Mackie 1970	Dunak 1971	Jorgens. 1970	Harvey 1982	Cul/Boov 1982	Culwell 1986	Progers 1978	Hansen 1985	Mus./Sta. 1980	MTNHP
.	.	.	.	.	.	.	.	.	.	Foid/Agca	Foid-Agca
.	.	.	.	.	.	.	.	.	.	Foid/Agca	Foid-Agca
.	.	.	.	.	.	.	.	.	.	Foid/Agcp	Foid-Agcp
.	.	.	.	.	.	.	.	.	Foid/Caha	.	Foid-Caha
.	.	.	.	.	.	.	.	.	.	Foid/Caac	Foid-Caac
.	.	.	.	.	.	.	.	.	.	Foid/Stri	Foid-Stri
.	.	.	.	.	.	.	.	.	.	Fpac/Agcp	Fpac-Agcp
.	.	.	.	.	.	.	.	.	.	Fpac/Foid	Fpac-Foid
.	.	.	Slips (in part)	.	Stoo/Bogr	Stoo/Agam	Stoo/Ca/V Bogr	Stoo/Bogr- Cafi	.	Stoo/Bogr	Stoo-Bogr
.	.	.	Slips (in part)	.	.	Stoo/Ca/V Kocr	.	.	Stoo/Cafi	.	Stoo-Cafi
.	.	.	.	.	.	.	.	.	Stoo/Caha	.	Stoo-Caha

<sup>1</sup> authors describing associations not described by the other authors listed.

<sup>2</sup> *Agropyron spicatum* community where *Slips comata* is often well-represented.

Table 2. Relationship of preliminary eastern Montana shrubland plant communities identified by the MTNHP to previous classifications. Riparian and tundra communities are not included.

other <sup>1</sup>	Brown 1971	Mackie 1970	Dusak 1971 Arca-Agro	Jorgens 1979 Arca/Agam	Harvey 1982 Arca/Agam	Cul/Scow 1982 Arca/Agam	Culwell 1986 Arca/Elam	Progers 1978 Arca/Agam/ Boqr	Hansen 1985 Arca/Agam	Mus./Sta. 1980	MTNHP Arca/Agam
.	.	.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.	Arca/Feld	Arca/Feld
.	.	.	.	.	.	.	.	Arca/Bivr- Agam	.	.	Arca/Bivr
.	.	Arlo	.	.	Arlo/Calc	.	.	.	.	.	Arlo
.	.	.	.	.	.	.	.	.	.	Arar/Agpp	Arca/Agpp
.	.	.	.	.	.	.	.	.	.	Arar/Feld	Arca/Feld
.	.	.	.	Artr/Agla, Agpp ph.	.	.	.	.	.	.	Artr/Agla
.	.	Artr-Agam	Artr-Agro	Artr/Agpp, Agam ph.	.	Artr/Agam	Artr/Elam	Artr/Agam/ Boqr	Artr/Agam	.	Artr/Agam
.	Artr/Agpp	Artr-Agpp	.	Artr/Agpp, Boqr ph.	Artr/Agpp	Artr/Agpp	.	.	Artr/Agpp	Artr/Agpp	Artr/Agpp
.	Alco-Artr, Artr-Alco- Agpp	.	.	.	.	+/- breaks complex	Artr/Alco; Alco/Artr	Artr bad- lands (in part)	.	.	Artr-Alco
.	.	.	.	.	Artr/Alnu	.	.	.	.	.	Artr-Alnu
.	.	.	.	Artr/Feld	.	.	.	.	.	.	Artr/Feld
.	.	.	.	.	.	.	.	.	.	Artr/Fesc	Artr/Fesc
.	.	.	.	.	Artr/Bleo	.	.	.	.	.	Artr/Bleo
.	.	.	.	.	Alnu/Agam	.	.	.	.	.	Alnu
.	.	.	.	.	.	.	.	.	.	Calc/Agpp	Calc/Agpp
.	.	.	.	.	.	Juho side- hill	.	Juho/Arac- Agpp	.	.	Juho/Arac
.	.	Juho	.	.	.	.	.	.	Juho/Calc	.	Juho/Calc
.	.	.	.	.	.	.	.	.	.	Poftr/Feld	Poftr/Feld
.	.	.	.	.	.	.	.	.	.	Poftr/Fesc	Poftr/Fesc

Table 2. (continued)

other	Brown 1971	Mackie 1970	Dusak 1971	Jorgens. 1979	Harvey 1982	Cul/Sow 1982	Culwell 1988	Progers 1978	Hansen 1985	Mus./Sis. 1980	MTNHP
.	.	.	.	.	.	.	.	.	.	Putr/Fold	Putr/Fold
.	.	.	.	.	.	.	.	.	.	Putr/Fanc	Putr/Fanc
.	.	.	decid. alrub	.	.	Rhr/Agpp	.	.	Rhr/Agpp	Rhr/Agpp	Rhr/Agpp
Cul. '85	.	.	.	.	.	.	.	.	.	.	Rhr/Anas
.	.	.	.	.	.	.	.	.	Rhr/Cafi	.	Rhr/Cafi
.	.	.	.	.	.	Rhr/Sloc; Rhr/side- hill	.	.	.	.	Rhr/Cabo
.	.	.	.	.	.	.	.	.	Rhr/Fold	Rhr/Fold	Rhr/Fold
.	.	.	.	Beve/Agda	.	.	.	.	.	.	Beve/Agda
.	.	Beve-Agum (in part)	.	.	Beve/Agum; Beve/Alnu	.	.	.	Beve/Agum	Beve/Agum	Beve/Agum
.	Beve	.	.	.	.	.	.	.	Beve/Agpp	.	Beve/Agpp
.	.	Beve-Agum (in part)	Artir-Beve; Beve	Artir/Agda, Beve ph.	Beve/Artir; Beve/Artir- Alnu	.	Beve	.	.	.	Beve/Artir

1 authors describing associations not described by the other authors listed.

\* \* D R A F T \* \*  
August 27, 1990

- a) predominant surface geology map unit
- b) predominant soil map unit
- c) altitude (at center of grid cell)
- d) estimated mean annual rainfall
- e) estimated solar radiation

Grid cells representing the range of these characteristics within BLM lands will be selected for field sampling to ensure representation of the total environmental variation present. Each sample will consist of one to five plots chosen from different topographic positions and where subjective judgement indicates a marked change in vegetation composition.

To minimize the confounding nature of heavy disturbance on vegetation occurrence, areas severely overgrazed, herbicide treated, mechanically disturbed, artificially seeded, or irrigated will not be sampled. Plots will be established within portions of stands that appear to be relatively uniform in topography and vegetation structure.

Complete taxonomic lists of vascular plants will be recorded within each 375 m<sup>2</sup> circular study plot. Canopy cover classes of each species will be estimated. The data gathered will include completing a Natural Heritage Program Community Survey Form for each plot.

Analysis will focus on using a combination of classification to determine community types, and gradient analyses to describe general patterns of communities/species in relation to environmental factors. Classification will initially

be achieved using the objective TWo-way INDicator SPecies ANalysis (TWINSPAN: Hill 1979a) procedure and cluster analysis (CA: Lance and Williams 1967). The final classification will be based on a combination of the objective analyses, subjective ecological judgement, and consideration of the gradient analysis results.

The gradient analyses will utilize a combination of objective procedures, including DETrended CORrespondence ANALysis (DECORANA: Hill 1979b) and Nonmetric MultiDimensional Scaling (NMDS: Ludwig and Reynolds 1988), and subjective graphic descriptions of community/environmental patterns (Whittaker 1967). CANOnical CORrespondence analysis (CANOCO: Ter Braak 1986) and Multiple Discriminant Analysis (MDA: Ludwig and Reynolds 1988) will be used to statistically test the degree the species and communities are related to supplied environmental variables.

### OUTPUTS

	<u>Date</u>	<u>Content</u>
A	05/91	Preliminary classification of grassland/shrubland communities on BLM lands in Montana based on existing data and literature.
B	06/91	Descriptions and maps of grid cells selected for sampling in 1991.
C	12/91	Refined preliminary classification of Montana grassland and shrubland communities on BLM lands based on 1991 fieldwork.

\* \* D R A F T \* \*  
August 27, 1990

- D     06/92     Descriptions and maps of grid cells selected for sampling in 1992.
  
- E     04/93     "Final" classification of Montana grassland and shrubland communities on BLM lands including discussion of sensitivities and potential responses to perturbation, and identification of high priority communities and sites for conservation.
  
- F     07/93     Following critical review of Output E by the BLM, the classification will be published. Also, a journal article will be submitted that describes grassland-shrubland community /environmental patterns in Montana.

*CURRENT AND POTENTIAL SUPPORT*

The Nature Conservancy currently provides salary and limited field-work support for R. DeVelice. A project of the magnitude proposed will require additional personnel for field and office assistance and an expanded travel budget.

\* \* D R A F T \* \*  
August 27, 1990

*ESTIMATED BUDGET*

Project Duration: April 1, 1991 - April 30, 1993

Direct Costs

Salaries and Wages

Robert L. DeVelice, Ph.D.

1991-92, 12 mos., 0.1 FTE      \$ 3000

1992-93, 12 mos., 0.1 FTE      3500

field assistant

1991, 3 mos., 1.0 FTE      5000

1992, 3 mos., 1.0 FTE      5500

office assistant

1991, 1 mo., 1.0 FTE      1000

1992, 1 mo., 1.0 FTE      1000

Total Salaries and Wages      \$19000

Fringe Benefits (approx. 15% Sal. & Wages)      \$ 2500

Materials and Supplies

topographic maps      \$ 300

miscellaneous      500

Total Materials and Supplies      \$ 800

\* \* D R A F T \* \*  
August 27, 1990

Travel

fieldwork (5000 mis. @ \$0.27/mi.)	\$1350
"Per diem" (20 days @ \$25/day)	500
Total Travel	\$ 1850
Total Direct Costs (TDC)	\$24150
Total Indirect Costs (circa 30% of TDC)	\$ 5850
<b>TOTAL PROJECT COSTS</b>	<b><u>\$30,000</u></b>

\* \* D R A F T \* \*  
August 27, 1990

*LITERATURE CITED*

- Anderson, N.L. 1973. The vegetation of rangeland sites associated with some grasshopper studies in Montana. Montana Agriculture Experiment Station Bulletin No. 668. Montana State University, Bozeman.
- Austin, M.P. and P.C. Heyligers. 1989. Vegetation survey design for conservation: gradsect sampling of forests in north-eastern New South Wales. *Biological Conservation* 50:13-32.
- Boggs, K.W. 1984. Succession in riparian communities of the lower Yellowstone River, Montana. Unpublished M.S. Thesis, Montana State University, Bozeman.
- Brown, R.W. 1971. Distribution of plant communities in southeastern Montana badlands. *American Midland Naturalist* 85:458-477.
- Cooper, S.V. and R.D. Pfister. 1981. Forest habitat types of the Blackfeet Indian Reservation. Review draft, 5/21/81, for Bureau of Indian Affairs, Wind River Agency, Fort Washakie, Wyoming.
- Cooper, S.V. and R.D. Pfister. 1985. Forest habitat types of the Crow and Northern Cheyenne Indian Reservations. Termination report to Bureau of Indian Affairs, Billings Area Office (draft). USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah.
- Coupland, R.T. 1950. Ecology of mixed prairie in Canada. *Ecological Monographs*. 20:271-315.
- Coupland, R.T. 1961. A reconsideration of grassland classification in the northern Great Plains of North America. *Journal of Ecology* 49:135-167.
- Culwell, L.D. and Scow, K.L. 1982. Terrestrial vegetation inventory: Dominy project area, Custer County, MT 1979-1980. Unpublished Technical Report for Western Energy Company by Westech. Helena, Montana.
- Culwell, L.D., Scow, K.L., and Larsen, L.A. 1985. Vegetation baseline inventory, Peabody Coal Company, Big Sky Mine Area B, Rosebud County, Montana. Westech. Helena, Montana.
- Culwell, L.D., Scow, K.L., and Larsen, L.A. 1986. A preliminary evaluation of the vegetation of the Makoshika State Park area, Dawson County, Montana (draft). Unpublished report for Montana Department of Fish, Wildlife and Parks. Westech, Helena, Montana.

\* \* D R A F T \* \*  
August 27, 1990

- Dusek, G.L. 1971. Range relationships of mule deer in the prairie habitat, northcentral Montana. Unpublished M.S. thesis, Montana State University, Bozeman.
- Gillison, A.N. and K.R.W. Brewer. 1985. The use of gradient directed transects or gradsects in natural resource surveys. *Journal of Environmental Management* 20:103-127.
- Girard, M.M., H. Goetz, and A.J. Bjugstad. 1989. Native woodland habitat types of southwestern North Dakota. USDA Forest Service Research Paper RM-281.
- Hansen, P.L. 1985. An ecological study of the vegetation of the Grand River/Cedar River, Sioux, and Ashland Districts of the Custer National Forest, Ph.D., South Dakota State University.
- Hansen, P.L. and G.R. Hoffman. 1985. An ecological study of the vegetation of the Grand River/Cedar River, Sioux, and Ashland districts of the Custer National Forest: a habitat type classification. USDA Forest Service Cooperative Agreement No. RM-80-131-CA.
- Hansen, P.L., G.R. Hoffman, and A.J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: a habitat type classification. USDA Forest Service General Technical Report RM-113.
- Harvey, S.J. 1982. Vegetation of Musselshell and Petroleum Counties, Montana. Unpublished report to the Soil Conservation Service, Roundup, Montana.
- Hill, M.O. 1979a. TWINSpan: A FORTRAN program for arranging multivariate data in an ordered two-way table by classification of the individuals and attributes. *Ecology and Systematics*, Cornell University, Ithaca, New York.
- Hill, M.O. 1979b. DECORANA: A FORTRAN program for detrended correspondence analysis and reciprocal averaging. *Ecology and Systematics*, Cornell University, Ithaca, New York.
- Hoffman, G.R. and P.L. Hansen. 1981. Habitat types of the Custer National Forest. USDA Forest Service Contract No. RM-RFP-80-131-CA.
- Jones, G. 1989. Preliminary Wyoming plant community classification (draft). Wyoming Natural Heritage Program, Laramie.
- Jorgensen, H.E. 1979. Vegetation of the Yellow Water Triangle, Montana. Montana Department of Fish and Game, in cooperation with the Bureau of Land Management. Helena, Montana.

\* \* D R A F T \* \*  
August 27, 1990

- Lance, G.N. and W.T. Williams. 1967. A general theory for classificatory sorting strategies. 1. hierarchical systems. *Computer Journal* 10:271-276.
- Ludwig, J.A. and J.F. Reynolds. 1988. *Statistical ecology: a primer on methods and computing*. John Wiley & Sons, New York.
- Mackie, R.J. 1970. Range ecology and relations of mule deer, elk, and cattle in the Missouri River breaks, Montana. *Wildlife Monographs* 20:1-79.
- MTNHP. 1990. Guide to the natural vegetation of Montana (draft). Montana Natural Heritage Program, Helena.
- Mueggler, W.F. and W.L. Stewart. 1980. Grassland and shrubland habitat types of western Montana. USDA Forest Service General Technical Report INT-66.
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. *Annals of the Association of American Geographers* 77:118-125.
- Pfister, R.D., B.L. Kovalchik, S.F. Arno, and R.C. Presby. 1977. Forest habitat types of Montana. USDA Forest Service General Technical Report INT-34.
- Progers, R. 1978. Circle West vegetation baseline study: final report. Circle West Technical Report No. 1. Energy Division, Montana Department of Natural Resources and Conservation, Helena.
- Roberts, D.W. 1980. Forest habitat types of the Bear's Paw Mountains and Little Rocky Mountains, Montana. Unpublished M.S. thesis, University of Montana, Missoula.
- Ross, R.L., Murray, E.P., and Haigh, J.G. 1973. Soil and vegetation inventory of near-pristine sites, Montana. USDA Soil Conservation Service, Bozeman, Montana.
- Taylor, J.E. and Holst, T.L. 1976. Grass and shrub plant community classification. USDA Forest Service contract research project 26-3494, final report, Ashland District, USDA Forest Service.
- Ter Braak, C.J.F. 1986. Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology* 67:1167-1179.
- Whitman, W. and H.C. Hanson. 1939. Vegetation on scoria and clay buttes in western North Dakota. *Ecology* 20:455-457.
- Whittaker, R.H. 1967. Gradient analysis of vegetation. *Biological Review* 42:207-264.