

# *Lactuca serriola* (Prickly Lettuce)

## Predicted Suitable Habitat Modeling

**Distribution Status:** Present

**State Rank:** [SNA](#) (Non-native Species)

**Global Rank:** [GNR](#)

### Modeling Overview

**Created by:** Braden Burkholder

**Creation Date:** April 27, 2023

**Evaluator:** Bryce Maxell

**Evaluation Date:** May 3, 2023



**Inductive Model Goal:** To predict the current distribution and relative suitability of general habitat for *Lactuca serriola* at large spatial scales across its presumed current range in Montana.

**Inductive Model Performance:** The model appears to generally reflect the current distribution and relative suitability of general habitat for *Lactuca serriola* at larger spatial scales across its presumed current range in Montana.

Evaluation metrics indicate a good model fit and the delineation of habitat suitability classes is well supported by the data. The model is presented as a reference, but more observation records, site-specific data, and/or other environmental layers may be needed to improve performance.

**Inductive Model Output:** [http://mtnhp.org/models/files/Lactuca\\_serriola\\_PDAST5F090\\_20230427\\_modelHex.lpk](http://mtnhp.org/models/files/Lactuca_serriola_PDAST5F090_20230427_modelHex.lpk)

**Suggested Citation:** Montana Natural Heritage Program. 2023. *Lactuca serriola* (Prickly Lettuce) predicted suitable habitat model created on April 27, 2023. Montana Natural Heritage Program, Helena, MT. 17 pp.

**Montana Field Guide Species Account:** <http://fieldguide.mt.gov/speciesDetail.aspx?elcode=PDAST5F090>

**Species Model Page:** <http://mtnhp.org/models/?elcode=PDAST5F090>

## **Inductive Modeling**

### **Model Limitations and Suggested Uses**

This model is based on statewide biotic and abiotic environmental layers originally mapped at a variety of spatial scales and standardized to 90×90-meter raster pixels. The spatial accuracy of the training and testing data are varied (typically 20-400 meters) and may result in additional statistical noise in the model. As a result, model outputs may not be appropriate for use on smaller areas or at fine spatial scales. **Model outputs should not typically be used for planning efforts on land areas smaller than one quarter of a public land survey system (PLSS) section (<64 hectares)** and model outputs for some species may only be appropriate for broader regional level planning efforts. Models should be interpreted as landscape-level habitat suitability (fundamental niche) and not as estimated distributions of the species (realized niche) since suitable habitat may be unoccupied (Pulliam 2000). **Consequently, model outputs should not be used in place of on-the-ground surveys for species**, and wildlife and land management agency biologists should be consulted about the value of using model output to guide habitat management decisions for regional planning efforts or local projects. See Suggested Contacts for Natural Resource Agencies listed at the end of this report or on [our website](#). In general, we have found across a large number of species representing a wide variety of plant and animal taxa that experts believe optimal and moderate suitability classes represent landscapes where suitable habitat is often more continuous while the low suitability class represents landscapes where suitable habitat is often less continuous, scattered, or patchy (see definitions in the Model Outputs and Evaluation section below). We encourage use of these classes for management, planning, permitting, survey, and other decisions accordingly.

## **Inductive Model Methods**

### **Modeling Process**

Presence-only data were extracted from Montana Natural Heritage Program Databases, which serve as a clearinghouse for animal and plant observation data in Montana. These data were then filtered to ensure spatial and temporal accuracy and to reduce spatial autocorrelation (summarized in Table 1). The spatial extent of this model was limited to the presumed geographic range of the species, by season when applicable, in order to accurately assess potentially available habitat.

We then used these data and 44 statewide biotic and abiotic environmental layers at a 90×90-meter pixel scale (Table 2) to construct the model using a maximum entropy algorithm employed in the modeling program Maxent (Phillips et al. 2006, Phillips et al. 2017). Entropy maximization modeling functions by calculating constraint distributions and then applies those constraints to the environmental layers to estimate a predicted suitable habitat distribution. The mean and variance of the environmental layer values (environmental variables) at the training data locations are used to estimate the constraint distributions. Maxent requires that the final predicted distribution fulfills these constraints. Maxent avoids overfitting models to the training data by “regularizing” or relaxing the constraints so that modeled distributions only have to be close to, rather than exactly equal to, the constraint distributions (Elith et al. 2011). The default regularization multiplier of 1.0 was used since species-specific tuning was impractical given the diversity and volume of species modeled in this effort (Merow et al. 2013, Radosavljevic and Anderson 2014). Additionally, we did not use hinge or threshold features at any sample size to minimize potential overfitting by overly complex models (Syfert et al. 2013, De Marco and Nóbrega 2018). The Maxent algorithm can successfully train models even when collinearity exists between environmental variables and the practices of removing collinear variables and/or reducing variables results in limited improvement in Maxent model performance (De Marco and Nóbrega 2018, Feng et al. 2019); neither method was employed here.

Maxent fits a model by assuming the predicted distribution is uniform in geographic space and moves away from this distribution only to the extent that it is forced to by the constraints of the training data. To do this, Maxent successively modifies the coefficients for each environmental variable via random walk, accepting the modified coefficient if it increases the gain. Gain is a measure of the closeness of the model concentration around the presence samples that is similar to goodness of fit in generalized linear models. The random walk of coefficients continues until either the increase in the gain falls below a set convergence threshold (0.00001) or a set maximum number of iterations are performed (50,000). The gain value at the end of a model run indicates the likelihood of suitability of the presence samples relative to the likelihood for random background points.

We employed a k-folds cross validation methodology, in this case using ten folds for model training and validation (Elith et al. 2011). Each fold consists of 90% of the data designated for training and 10% of the data reserved for testing. Each record is used for training nine times and testing once. Ten models are estimated and averaged to produce the final model presented here.

### Model Outputs and Evaluation

The overall gain associated with individual environmental variables (Table 3) can be used as a measure of the relative importance of each variable (Merow et al. 2013). However, the importance of individual environmental variables should be interpreted with caution due to collinearity between variables. The jackknife assessment of contribution by individual environmental variables to training gain (Figure 1) may be more useful in interpreting the relative importance of individual variables. The response curves for the top four contributing environmental variables are shown for reference (Figure 2). These response curves should also be interpreted cautiously because the observation data used to train the models was not gathered under a probabilistic sampling scheme. If enough observations were available to train and evaluate the model, thresholds are estimated for low, moderate, and optimal habitat suitability; details of this process are presented in Table 4 and Figure 3.

The initial model output is a spatial dataset of continuous logistic values that ranges from 0-1 with lower values representing areas predicted to be less suitable habitat and higher values representing areas predicted to be more suitable habitat (Figures 4 & 5). The standard deviation in the model output across the averaged models is also calculated and plotted as a map to examine spatial variance of model output (Figure 6). The continuous output is reclassified into suitability classes and aggregated within 259-hectare hexagons (Figures 7-9).

In addition to the map of spatial variance in model output, we also evaluated the output of the Maxent model with absolute validation index (AVI) (Hirzel et al. 2006) and deviance (Phillips and Dudik 2008). These metrics are described below in the results (Table 5). True skill statistic (TSS) (Allouche et al. 2006), symmetric extremal dependence index (SEDI) (Wunderlich et al. 2019), and Area Under the Curve (AUC) values are also displayed for reference but are not used for evaluation (Lobo et al. 2008). Finally, a deviance value was calculated for each test data observation as a measure of how well model output matched what the model predicted for the location of test observations and this was plotted with larger symbols indicating larger deviance (see Figure 5). In practice, we have found large deviance values to be associated with records that are incorrectly or imprecisely mapped, problematic areas in underlying environmental layers, regions where species have few observations outside of the core portion of their range, or insufficient models with poor performance.

Table 1: Model Data Selection Criteria and Summary

<b>Location Data Source</b>	Montana Natural Heritage Program Databases
<b>Total Number of Records</b>	722
<b>Location Data Selection Rule 1 – Valid and Accurate Records</b>	Records with <= 800 meters of locational uncertainty for years after 1960
<b>Number of Locations Meeting Selection Rule 1</b>	666
<b>Location Data Selection Rule 2 – Spatially Unique Records</b>	No overlap in locations within 3200 meters in order to avoid spatial autocorrelation
<b>Observation Records used in Model (Locations Meeting Selection Rules 1 &amp; 2)</b>	498
<b>Number of Model Background Locations<sup>a</sup></b>	60,000

<sup>a</sup> Background locations are chosen at random and in proportion to the percent of the state covered by a species' geographic range, with a maximum of 60,000 locations. Although these locations only represent ~0.1% of the pixels in any modeled area, this level of sampling is sufficient to estimated distributions of environmental conditions present (Phillips and Dudik 2008).

Table 2: Environmental Layers and Corresponding Variables<sup>a</sup>

Layer Name	Variable	Layer Name	Variable
LC_AgDry_97	Developed - Dry Agriculture	NED_AspectEW	Aspect (East-West)
LC_AgIrr_97	Developed - Irrigated Lands	NED_AspectNS	Aspect (North-South)
LC_Alpine_97	Alpine	NED_Elevation	Elevation
LC_Barren_97	Sparse and Barren	NED_Ruggedness	Ruggedness
LC_Developed_97	Developed - All Other	NED_Slope	Slope
LC_ForestBurn_97	Forest - Burned	NED_SRISummer	Summer Solar Radiation
LC_ForestConif_97	Forest - Conifer	NED_SRIWinter	Winter Solar Radiation
LC_ForestDecid_97	Forest - Deciduous	NED_TPI	Topographic Position Index
LC_ForestHarv_97	Forest - Harvested	NHD_Dist2WaterEdge	Distance to Water Edge
LC_ForestInsct_97	Forest - Insect Killed	NHP_AnthroInfl	Anthropogenic Influence
LC_Grassland_97	Grasslands	NRCS_FrostFreeDays	Frost Free Days
LC_IntroVeg_97	Introduced Vegetation	NRCS_REAP	Relative Effective Annual Precipitation
LC_ShrubBurn_97	Shrublands - Burned	PRISM_Precipitation	Annual Precipitation
LC_Shrubland_97	Shrublands	PRISM_WinPrecip	Percent Winter Precipitation
LC_WetRip_97	Wetland & Riparian	SoilGrid_BD	Bulk Density
LC_Dist2Forest	Distance to Forest	SoilGrid_Clay	Percent Clay
MCO_DegreeDays	Degree Days	SoilGrid_EC	Electric Conductivity
MCO_MaxSumTemp	Maximum Summer Temp	SoilGrid_OrgC	Organic Carbon
MCO_MinWinTemp	Minimum Winter Temp	SoilGrid_pH	Soil pH
MCO_NDVI	Normalized Difference Vegetation Index	SoilGrid_Sand	Percent Sand
MTGeol_Dist2Alluv	Distance to Alluvium	SoilGrid_Silt	Percent Silt
MTGeol_Dist2C03	Distance to Carbonate Rock	SoilGrid_TotN	Total Nitrogen

<sup>a</sup> Additional details and sources available in Appendix.

## Inductive Model Results

Table 3: Top Ten Contributing Environmental Variables to Model Fit

Variable	Percent Contribution <sup>a</sup>	Variable	Percent Contribution <sup>a</sup>
% Winter Precipitation	15.5%	Summer Solar Radiation	5.2%
Frost Free Days	10.8%	Bulk Density	4.4%
Degree Days	10.1%	Forest - Conifer	4.0%
Developed - Dry Agriculture	6.9%	Soil pH	3.7%
Maximum Summer Temp	6.9%	Grasslands	3.3%

<sup>a</sup> Relative contributions of the variables to the model based on changes in fit (gain) during iterations of the training algorithm.

Table 4: Habitat Suitability Thresholds and Areas of Suitable Habitat

Measure	Value
Optimal Logistic Threshold <sup>a</sup>	0.831
<b>Moderate Logistic Threshold<sup>b</sup></b>	<b>0.210</b>
Low Logistic Threshold <sup>c</sup>	0.054
Area of predicted optimal habitat within modeled range	918.8 km <sup>2</sup>
<b>Area of predicted moderate suitability habitat within modeled range</b>	<b>125,258.9 km<sup>2</sup></b>
Area of predicted low suitability habitat within modeled range	84,941.0 km <sup>2</sup>
Total area of predicted suitable habitat within modeled range	211,118.7 km <sup>2</sup>
Area of entire modeled range (percent of Montana)	380,529.0 km <sup>2</sup> (100.0%)

<sup>a</sup> The logistic threshold where the percentage of test observations above the threshold is 10 or more times higher than would be expected if the observations were randomly distributed across logistic value classes (Hirzel et al. 2006) (see Figure 3). When sample sizes are small, it may be undetermined.

<sup>b</sup> **This is the cutoff recommended for use in management decisions.** The logistic threshold value where the percentage of test observations above the threshold is greater than what would be expected if the observations were randomly distributed across logistic value classes - in other words, when the modeled habitat is used more often than expected from its proportional availability on the landscape (Hirzel et al. 2006). When sample sizes are small, it may be undetermined.

<sup>c</sup> The logistic threshold between unsuitable and low suitability as determined by Maxent which balances data omission error with minimizing predicted suitable area (Phillips et al. 2017). This is a conservative threshold that should encompass nearly all potentially suitable habitat for a species. In practice, habitat with low suitability may represent landscapes of marginal or discontinuous habitat where suitable habitat patches of various sizes are isolated by unsuitable habitat.

Table 5: Evaluation Metrics

Metric	Value
Low AVI <sup>a</sup>	95.8%
Moderate AVI <sup>a</sup>	81.5%
Optimal AVI <sup>a</sup>	4.0%
Average Testing Deviance ( $\bar{x} \pm sd$ ) <sup>b</sup>	2.037 $\pm$ 1.714
TSS (Sensitivity + Specificity - 1) <sup>c</sup>	0.4887 (0.8153 + 0.6735 - 1)
SEDI <sup>c</sup>	0.6481
Training AUC <sup>d</sup>	0.870
Test AUC <sup>e</sup>	0.824

<sup>a</sup> Absolute Validation Index: The proportion of test locations that fall above the low, moderate, or optimal logistic threshold (see Table 4).


<sup>b</sup> A measure of how well model output matched the location of test observations. In theory, everywhere a test location was located, the logistic value should have been 1.0. The deviance value for each test location is calculated as -2 times the natural log of the associated logistic output value. For example, the equivalent deviance values for the low, moderate and optimal logistic thresholds of this model would be 5.841, 3.117 and 0.371, respectively. Deviances for individual test locations are plotted in Figure 5. Average Testing Deviance less than the Moderate Deviance typically indicates good model performance.

<sup>c</sup> Ranges from -1 to 1, with a random or null model performing at a value of 0 and values >0.65 indicating moderate performance (>0.8 generally good performance). The moderate threshold (0.210) is used to develop the confusion matrix for Sensitivity and Specificity metrics. Note that Specificity is calculated based on pseudo-absences (not true absences) and may be biased when large areas are modeled as moderate or optimal suitable habitat.

<sup>d</sup> The area under a curve obtained by plotting the true positive rate against 1 minus the false positive rate for model training observations (averaged over 10 folds). Values range from 0 to 1 with a random or null model performing at a value of 0.5.

<sup>e</sup> The same metric described in <sup>d</sup>, but calculated for test observations.

Figure 1. Jackknife assessment of contribution by individual environmental variables to training gain. Variables are ordered by reduction in gain without that variable (green), from greatest to least impact. Only the 25 most influential variables<sup>a</sup> are shown.



<sup>a</sup> Interpretation of individual environmental variables should be approached cautiously and may be inappropriate due to covariance between variables.

Figure 2. Response curves for the top four contributing environmental variables, mean value in red, +/- one standard deviation in blue. Response curves for additional environmental variables are available upon request.





Figure 3. Thresholds for moderate and optimal suitability classes as determined by logarithmic fit.





## Inductive Model Map Outputs

Figure 4. Continuous habitat suitability model logistic output (90-meter pixels); white area is not modeled.




Figure 5. Continuous habitat suitability model output with relative deviance for each observation. Low deviance points fall within optimal or moderate habitat; high deviance points are in generally unsuitable habitat.






Figure 6. Standard deviation in the model output across the averaged models. Lower deviance (a solid blue map) indicates a better fitting model with lower variability between model iterations.




Figure 7. Model output for 90-meter pixels classified into habitat suitability classes.






Figure 8. Model output classified into habitat suitability classes and aggregated into hexagons at a scale of 259 hectares per hexagon. This is the finest scale suggested for management decisions and survey planning.





Figure 9. Model output classified into habitat suitability classes and aggregated into hexagons; observations used for modeling are displayed for reference.



## Literature Cited

- Allouche, O., A. Tsoar, and R. Kadmon. 2006. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology* 43: 1223-1232. <https://doi.org/10.1111/j.1365-2664.2006.01214.x>
- De Marco, P. Jr. and C. C. Nóbrega. 2018. Evaluating collinearity effects on species distribution models: An approach based on virtual species simulation. *PLoS ONE* 13(9): e0202403. <https://doi.org/10.1371/journal.pone.0202403>
- Elith, J., S. J. Phillips, T. Hastie, M. Dudík, Y. E. Chee, and C. J. Yates. 2011. A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions* 17: 43-57. <https://doi.org/10.1111/j.1472-4642.2010.00725.x>
- Feng, X., D. S. Park, Y. Liang, R. Pandey, M. Papeş. 2019. Collinearity in ecological niche modeling: Confusions and challenges. *Ecology and Evolution* 9: 10365-10376. <https://doi.org/10.1002/ece3.5555>
- Hirzel, A. H., G. Le Lay, V. Helfer, C. Randin, and A. Guisana. 2006. Evaluating the ability of habitat suitability models to predict species presences. *Ecological Modelling* 199: 142-152. <https://doi.org/10.1016/j.ecolmodel.2006.05.017>
- Lobo, J. M., A. Jiménez-Valverde, and R. Real. 2008. AUC: a misleading measure of the performance of predictive distribution models. *Global Ecology and Biogeography* 17: 145-151. <https://doi.org/10.1111/j.1466-8238.2007.00358.x>
- Merow, C., M. J. Smith, and J. A. Silander Jr. 2013. A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography* 36: 1058-1069. <https://doi.org/10.1111/j.1600-0587.2013.07872.x>
- Phillips, S. J., R. P. Anderson, and R. E. Schapire. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modeling* 190: 231-259. <https://doi.org/10.1016/j.ecolmodel.2005.03.026>
- Phillips, S. J., M. Dudík, R. E. Schapire. 2017. Maxent software for modeling species niches and distributions (Version 3.4.1). Available from: [http://biodiversityinformatics.amnh.org/open\\_source/maxent](http://biodiversityinformatics.amnh.org/open_source/maxent)
- Phillips, S. J. and M. Dudík. 2008. Modeling of species distributions with MaxEnt: New extensions and a comprehensive evaluation. *Ecography* 31: 161-175. <https://doi.org/10.1111/j.0906-7590.2008.5203.x>
- Pulliam, H. R. 2000. On the relationship between niche and distribution. *Ecology Letters* 3: 349-361. <https://doi.org/10.1046/j.1461-0248.2000.00143.x>
- Radosavljevic, A., and R. P. Anderson. 2014. Making better Maxent models of species distributions: complexity, overfitting, and evaluation. *Journal of Biogeography* 41: 629-643. <https://doi.org/10.1111/jbi.12227>
- Syfert, M. M., M. J. Smith, and D. A. Coomes. 2013. The effects of sampling bias and model complexity on the predictive performance of Maxent species distribution models. *PLoS ONE* 8(2): e55158. <https://doi.org/10.1371/journal.pone.0055158>
- Wunderlich, R. F., Y. Lin, J. Anthony, and J. R. Petway. 2019. Two alternative evaluation metrics to replace the true skill statistic in the assessment of species distribution. *Nature Conservation* 35: 97-116. <https://doi.org/10.3897/natureconservation.35.33918>

## Appendix

Table A. Detailed Descriptions of Environmental Layers

Layer Name	Class	Variable	Original Scale	Description
LC_AgDry_97	Landcover	Developed - Dry Agriculture	30m	Proportion (0-1) of cells classed as Agriculture without Irrigation within a 97-cell neighborhood (~150m radius).
LC_AgIrr_97	Landcover	Developed - Irrigated Lands	30m	Proportion (0-1) of cells classed as Agriculture with Irrigation within a 97-cell neighborhood (~150m radius).
LC_Alpine_97	Landcover	Alpine	30m	Proportion (0-1) of cells classed as Alpine cover types within a 97-cell neighborhood (~150m radius).
LC_Barren_97	Landcover	Sparse and Barren	30m	Proportion (0-1) of cells classed as Sparse or Barren within a 97-cell neighborhood (~150m radius).
LC_Developed_97	Landcover	Developed - All Other	30m	Proportion (0-1) of cells classed as Developed (e.g. towns, roads, mines) within a 97-cell neighborhood (~150m radius).
LC_ForestBurn_97	Landcover	Forest - Burned	30m	Proportion (0-1) of cells classed as Burned Forest within a 97-cell neighborhood (~150m radius).
LC_ForestConif_97	Landcover	Forest - Conifer	30m	Proportion (0-1) of cells classed as Conifer Forest within a 97-cell neighborhood (~150m radius).
LC_ForestDecid_97	Landcover	Forest - Deciduous	30m	Proportion (0-1) of cells classed as Deciduous Forest within a 97-cell neighborhood (~150m radius).
LC_ForestHarv_97	Landcover	Forest - Harvested	30m	Proportion (0-1) of cells classed as Harvest Forest within a 97-cell neighborhood (~150m radius).
LC_ForestInsct_97	Landcover	Forest - Insect Killed	30m	Proportion (0-1) of cells classed as Insect Killed Forest within a 97-cell neighborhood (~150m radius).
LC_Grassland_97	Landcover	Grasslands	30m	Proportion (0-1) of cells classed as Grassland cover types within a 97-cell neighborhood (~150m radius).
LC_IntroVeg_97	Landcover	Introduced Vegetation	30m	Proportion (0-1) of cells classed as Introduced Vegetation within a 97-cell neighborhood (~150m radius).
LC_ShrubBurn_97	Landcover	Shrublands - Burned	30m	Proportion (0-1) of cells classed as Burned Shrubland within a 97-cell neighborhood (~150m radius).
LC_Shrubland_97	Landcover	Shrublands	30m	Proportion (0-1) of cells classed as Shrubland cover types within a 97-cell neighborhood (~150m radius).
LC_WetRip_97	Landcover	Wetland & Riparian	30m	Proportion (0-1) of cells classed as Wetland or Riparian within a 97-cell neighborhood (~150m radius).
LC_Dist2Forest	Landcover	Distance to Forest	30m	Distance in meters to any forest cover type, after consolidating to patches >900m <sup>2</sup> .
MCO_DegreeDays	Climate	Degree Days	800m	Average annual total of degree days (°F) above 32°F for 1981-2010.
MCO_MaxSumTemp	Climate	Maximum Summer Temp	800m	Average maximum temperature (°C) in July for 1981-2010.
MCO_MinWinTemp	Climate	Minimum Winter Temp	800m	Average minimum temperature (°C) in January for 1981-2010.
MCO_NDVI	Climate	Normalized Difference Vegetation Index	925m	Normalized Difference Vegetation as a measure of yearly mean greenness from the MODIS Terra satellite, 2000-2014.
MTGeol_Dist2Alluv	Geology	Distance to Alluvium	vector	Distance in meters to alluvial, glacial, or other unconsolidated surface geology types.
MTGeol_Dist2C03	Geology	Distance to Carbonate Rock	vector	Distance in meters to geological units with major components of either limestone or dolostone.



Layer Name	Class	Variable	Original Scale	Description
NED_AspectEW	Landform	Aspect (East-West)	≈10m	Aspect of slopes, ranging from 1 (east) to -1 (west).
NED_AspectNS	Landform	Aspect (North-South)	≈10m	Aspect of slopes, ranging from 1 (north) to -1 (south).
NED_Elevation	Landform	Elevation	≈10m	Elevation in meters above mean sea level.
NED_Ruggedness	Landform	Ruggedness	≈10m	Vector ruggedness measure from 0 (flat) to 1 (rugged), based on 5-cell neighborhood.
NED_Slope	Landform	Slope	≈10m	Percent slope (x100) of landscape.
NED_SRISummer	Landform	Summer Solar Radiation	≈10m	Solar radiation (WH/m <sup>2</sup> ) for the day of the summer solstice.
NED_SRIWinter	Landform	Winter Solar Radiation	≈10m	Solar radiation (WH/m <sup>2</sup> ) for the day of the winter solstice.
NED_TPI	Landform	Topographic Position Index	≈10m	Relative topographic position, based on a 5- to 10-cell radius envelope; positive values represent ridges, negative values are valleys/drainages.
NHD_Dist2WaterEdge	Hydrography	Distance to Water Edge	vector	Distance in meters to edges of lakes, reservoirs, streams, or rivers, as mapped in high resolution NHD.
NHP_AnthroInfl	Human	Anthropogenic Influence	vector	An index of human impact on landscape, based on structure density and road use.
NRCS_FrostFreeDays	Climate	Frost Free Days	30m	Estimated number of days without frost: daily minimum temperature > 0°C based on a 5 in 10 year probability.
NRCS_REAP	Climate	Relative Effective Annual Precipitation	10m	REAP, an adjusted annual precipitation estimate enhanced with DEM attributes (cm) over 30 years.
PRISM_Precipitation	Climate	Annual Precipitation	≈800m	Average annual precipitation (mm) for 1981-2010.
PRISM_WinPrecip	Climate	Percent Winter Precipitation	≈800m	Average percent (0 to 1) of the total annual precipitation that occurs during winter (Nov-Apr) for 1981-2010.
SoilGrid_BD	Soils	Bulk Density	100m	Bulk density (inversely related to pore space), g/cm <sup>3</sup> (0-5cm depth).
SoilGrid_Clay	Soils	Percent Clay	100m	Percent Clay in soil (0-5cm depth).
SoilGrid_EC	Soils	Electric Conductivity	100m	Electric conductivity (measure of salinity) of soil, dS/m (x100) (0-5cm depth).
SoilGrid_OrgC	Soils	Organic Carbon	100m	Soil organic carbon, % weight (x1000) (0-5cm depth).
SoilGrid_pH	Soils	Soil pH	100m	Soil pH (0-5cm depth).
SoilGrid_Sand	Soils	Percent Sand	100m	Percent Sand in soil (0-5cm depth).
SoilGrid_Silt	Soils	Percent Silt	100m	Percent Silt in soil (0-5cm depth); derived, based on Sand and Clay percentages.
SoilGrid_TotN	Soils	Total Nitrogen	100m	Total nitrogen in soil, % weight (x1000) (0-5cm depth).

#### Data Sources/Environmental Layer Name Prefix Key

LC - [Montana Land Use/Land Cover Dataset \(MSDI\)](#)

MCO - [Montana Climate Office \(MSDI\)](#)

MTGeol - [Montana Bureau of Mines & Geology \(MSDI\)](#)

NED - [National Elevation Dataset \(MSDI\)](#)

NHD - [National Hydrological Dataset \(MSDI\)](#)

NHP - [NHP Data \(unpublished\)](#)

NRCS - [Natural Resources Conservation Service \(MSDI\)](#)

PRISM - [PRISM Climate Group \(OSU\)](#)

SoilGrid - [US48 Soil Grids 100m \(PSU\)](#)



## Data Use Terms and Conditions


- Montana Natural Heritage Program (MTNHP) products and services are based on biological data and the objective interpretation of those data by professional scientists. MTNHP does not advocate any particular philosophy of natural resource protection, management, development, or public policy.
- MTNHP has no natural resource management or regulatory authority. Products, statements, and services from MTNHP are intended to inform parties as to the state of scientific knowledge about certain natural resources, and to further develop that knowledge. The information is not intended as natural resource management guidelines or prescriptions or a determination of environmental impacts. MTNHP recommends consultation with appropriate state, federal, and tribal resource management agencies and authorities in the area where your project is located.
- Information on the status and spatial distribution of biological resources produced by MTNHP are intended to inform parties of the state-wide status, known occurrence, or the likelihood of the presence of those resources. **These products are not intended to substitute for field-collected data, nor are they intended to be the sole basis for natural resource management decisions.**
- MTNHP does not portray its data as exhaustive or comprehensive inventories of rare species or biological communities. **Field verification of the absence or presence of sensitive species and biological communities will always be an important obligation of users of our data.**
- MTNHP responds equally to all requests for products and services, regardless of the purpose or identity of the requester.
- Because MTNHP constantly updates and revises its databases with new data and information, products will become outdated over time. Interested parties are encouraged to obtain the most current information possible from MTNHP, rather than using older products. We add, review, update, and delete records on a daily basis. Consequently, we strongly advise that you update your MTNHP data sets at a minimum of every three months for most applications of our information.
- MTNHP data require a certain degree of biological expertise for proper analysis, interpretation, and application. Our staff is available to advise you on questions regarding the interpretation or appropriate use of the data that we provide. Contact information for MTNHP staff is posted at: <http://mtnhp.org/contact.asp>
- The information provided to you by MTNHP may include sensitive data that if publicly released might jeopardize the welfare of threatened, endangered, or sensitive species or biological communities. This information is intended for distribution or use only within your department, agency, or business. Subcontractors may have access to the data during the course of any given project but should not be given a copy for their use on subsequent, unrelated work.
- MTNHP data are made freely available. **Duplication of hard-copy or digital MTNHP products with the intent to sell is prohibited without written consent by MTNHP.** Should you be asked by individuals outside your organization for the type of data that we provide, please refer them to MTNHP.
- MTNHP and appropriate staff members should be appropriately acknowledged as an information source in any third-party product involving MTNHP data, reports, papers, publications, or in maps that incorporate MTNHP graphic elements. The following is a suggested data citation format: *Montana Natural Heritage Program. {date type} for {species or species group} in {geographic filter, if applicable} Montana [Data set]. Retrieved January 1, 2020. Available from: <http://mtnhp.org>*
- Sources of our data include museum specimens, published and unpublished scientific literature, field surveys by state and federal agencies and private contractors, and reports from knowledgeable individuals. MTNHP actively solicits and encourages additions, corrections and updates, new observations or collections, and comments on any data we provide.
- MTNHP staff and contractors do not cross or survey privately-owned lands without express permission from the landowner. However, the program cannot guarantee that information provided to us by others was obtained under adherence to this policy.

## Suggested Contacts for Natural Resource Management Agencies


As required by Montana statute ([MCA 90-15](#)), the Montana Natural Heritage Program works with state, federal, tribal, nongovernmental organizations, and private partners to ensure that the latest animal and plant distribution and status information is incorporated into our databases so that it can be used to inform a variety of permitting and planning processes and management decisions. We encourage you to contact state, federal, and tribal resource management agencies in the area where your project is located and review the permitting overviews by the [Montana Department of Environmental Quality](#), the [Montana Department of Natural Resources and Conservation](#) and the [Index of Environmental Permits for Montana](#) for guidelines relevant to your efforts. In particular, we encourage you to contact the Montana Department of Fish, Wildlife, and Parks for the latest data and management information regarding hunted and high-profile management species and to use the U.S. Fish and Wildlife Service's [Information Planning and Consultation \(IPAC\) website](#) regarding U.S. Endangered Species Act listed Threatened, Endangered, or Candidate species.

For your convenience, we have compiled a list of relevant agency contacts and links below; check [our website](#) for updates.

### Montana Fish, Wildlife, and Parks

Fish Species	Zachary Shattuck <a href="mailto:zshattuck@mt.gov">zshattuck@mt.gov</a> (406) 444-1231 or Eric Roberts <a href="mailto:eroberts@mt.gov">eroberts@mt.gov</a> (406) 444-5334		
American Bison Black-footed Ferret Black-tailed Prairie Dog Bald Eagle Golden Eagle Common Loon Least Tern Piping Plover Whooping Crane	Kristina Smucker <a href="mailto:kismucker@mt.gov">kismucker@mt.gov</a> (406) 444-5209		
Grizzly Bear Greater Sage Grouse Trumpeter Swan Big Game Upland Game Birds Furbearers	Brian Wakeling <a href="mailto:Brian.Wakeling@mt.gov">Brian.Wakeling@mt.gov</a> (406) 444-3940		
Managed Terrestrial Game and Nongame Animal Data	Smith Wells – MFWP Data Analyst <a href="mailto:smith.wells@mt.gov">smith.wells@mt.gov</a> (406) 444-3759		
Fisheries Data	Ryan Alger – MFWP Data Analyst <a href="mailto:ryan.alger@mt.gov">ryan.alger@mt.gov</a> (406) 444-5365		
Wildlife and Fisheries Scientific Collector's Permits	<a href="https://fwp.mt.gov/buyandapply/commercialwildlifeandscientificpermits/scientific">https://fwp.mt.gov/buyandapply/commercialwildlifeandscientificpermits/scientific</a> Kammi McClain for Wildlife <a href="mailto:Kammi.McClain@mt.gov">Kammi.McClain@mt.gov</a> (406) 444-2612 Kim Wedde for Fisheries <a href="mailto:kim.wedde@mt.gov">kim.wedde@mt.gov</a> (406) 444-5594		
Fish and Wildlife Recommendations for Subdivision Development	Charlie Sperry <a href="mailto:CSperry@mt.gov">CSperry@mt.gov</a> (406) 444-3888 See also: <a href="https://fwp.mt.gov/conservation/living-with-wildlife/subdivision-recommendations">https://fwp.mt.gov/conservation/living-with-wildlife/subdivision-recommendations</a>		
Regional Contacts 	<a href="#">Region 1</a> - Kalispell (406) 752-5501 <a href="mailto:fwprg12@mt.gov">fwprg12@mt.gov</a> <a href="#">Region 2</a> - Missoula (406) 542-5500 <a href="mailto:fwprg22@mt.gov">fwprg22@mt.gov</a> <a href="#">Region 3</a> - Bozeman (406) 577-7900 <a href="mailto:fwprg3@mt.gov">fwprg3@mt.gov</a> <a href="#">Region 4</a> - Great Falls (406) 454-5840 <a href="mailto:fwprg42@mt.gov">fwprg42@mt.gov</a> <a href="#">Region 5</a> - Billings (406) 247-2940 <a href="mailto:fwprg52@mt.gov">fwprg52@mt.gov</a> <a href="#">Region 6</a> - Glasgow (406) 228-3700 <a href="mailto:fwprg62@mt.gov">fwprg62@mt.gov</a> <a href="#">Region 7</a> - Miles City (406) 234-0900 <a href="mailto:fwprg72@mt.gov">fwprg72@mt.gov</a>		

**Montana Department of Agriculture**General Contact Information: <https://agr.mt.gov/About/Office-Locations/Office-Locations-and-Field-Offices>Noxious Weeds: <https://agr.mt.gov/Noxious-Weeds>**Montana Department of Environmental Quality**Permitting and Operator Assistance for all Environmental Permits: <https://deq.mt.gov/Permitting>**Montana Department of Natural Resources and Conservation**Overview of, and contacts for, licenses and permits for state lands, water, and forested lands: <http://dnrc.mt.gov/licenses-and-permits>Stream Permitting (310 permits) and an overview of various water and stream related permits (e.g., Stream Protection Act 124, Federal Clean Water Act 404, Federal Rivers and Harbors Act Section 10, Short-term Water Quality Standard for Turbidity 318 Authorization, etc.). <http://dnrc.mt.gov/divisions/cardd/conservation-districts/the-310-law>Flood and Fire Resources: <http://dnrc.mt.gov/flood-and-fire>**Bureau of Land Management**

Montana Field Office Contacts:	
	
Billings	(406) 896-5013
Butte	(406) 533-7600
Dillon	(406) 683-8000
Glasgow	(406) 228-3750
Havre	(406) 262-2820
Lewistown	(406) 538-1900
Malta	(406) 654-5100
Miles City	(406) 233-2800
Missoula	(406) 329-3914


**United States Army Corps of Engineers**

Montana Regulatory Office for federal permits related to construction in water and wetlands

(406) 441-1375 <https://www.nwo.usace.army.mil/Missions/Regulatory-Program/Montana/>**United States Environmental Protection Agency**Environmental information, notices, permitting, and contacts <https://www.epa.gov/mt>Gateway to state resource locators <https://www.envcap.org/srl/index.php>**United States Fish and Wildlife Service**Information Planning and Conservation (IPAC) website: <https://ecos.fws.gov/ipac/>Montana Ecological Services Field Office (406) 449-5225 <https://www.fws.gov/montanafieldoffice/>**United States Forest Service**

Regional Office Contacts – Missoula			
Wildlife Program Leader	Tammy Fletcher	<a href="mailto:tammy.fletcher2@usda.gov">tammy.fletcher2@usda.gov</a>	(406) 329-3086
Wildlife Ecologist	Cara Staab	<a href="mailto:cara.staab@usda.gov">cara.staab@usda.gov</a>	(406) 329-3677
Fish Program Leader	Scott Spaulding	<a href="mailto:scott.spaulding@usda.gov">scott.spaulding@usda.gov</a>	(406) 329-3287
Fish Ecologist	Cameron Thomas	<a href="mailto:cameron.thomas@usda.gov">cameron.thomas@usda.gov</a>	(406) 329-3087
TES Program	Lydia Allen	<a href="mailto:lydia.allen@usda.gov">lydia.allen@usda.gov</a>	(406) 329-3558
Interagency Grizzly Bear Coordinator	Scott Jackson	<a href="mailto:scott.jackson@usda.gov">scott.jackson@usda.gov</a>	(406) 329-3664
Acting Regional Botanist	Amanda Hendrix	<a href="mailto:amanda.hendrix@usda.gov">amanda.hendrix@usda.gov</a>	(651) 447-3016
Regional Vegetation Ecologist	Mary Manning	<a href="mailto:marry.manning@usda.gov">marry.manning@usda.gov</a>	(406) 329-3304
Invasive Species Program Manager	Michelle Cox	<a href="mailto:michelle.cox2@usda.gov">michelle.cox2@usda.gov</a>	(406) 329-3669

## Tribal Nations



[Assiniboine & Gros Ventre Tribes – Fort Belknap Reservation](#)

[Assiniboine & Sioux Tribes – Fort Peck Reservation](#)

[Blackfeet Tribe - Blackfeet Reservation](#)

[Chippewa Creek Tribe - Rocky Boy's Reservation](#)

[Crow Tribe – Crow Reservation](#)

[Little Shell Chippewa Tribe](#)

[Northern Cheyenne Tribe – Northern Cheyenne Reservation](#)

[Salish & Kootenai Tribes - Flathead Reservation](#)

## Natural Heritage Programs and Conservation Data Centres in Surrounding States and Provinces

[Alberta Conservation Information Management System](#)

[British Columbia Conservation Data Centre](#)

[Idaho Natural Heritage Program](#)

[North Dakota Natural Heritage Program](#)

[Saskatchewan Conservation Data Centre](#)

[South Dakota Natural Heritage Program](#)

[Wyoming Natural Diversity Database](#)

## Invasive Species Management Contacts and Information

### Aquatic Invasive Species

[Montana Fish, Wildlife, and Parks Aquatic Invasive Species staff](#)

[Montana Department of Natural Resources and Conservation's Aquatic Invasive Species Grant Program](#)

[Montana Invasive Species Council \(MISC\)](#)

[Upper Columbia Conservation Commission \(UC3\)](#)

### Noxious Weeds

[Montana Weed Control Association Contacts Webpage](#)

[Montana Biological Weed Control Coordination Project](#)

[Montana Department of Agriculture - Noxious Weeds](#)

[Montana Weed Control Association](#)

[Montana Fish, Wildlife, and Parks - Noxious Weeds](#)

[Montana State University Integrated Pest Management Extension](#)

[Integrated Noxious Weed Management after Wildfires](#)

[Fire Management and Invasive Plants](#)