

Boreal Chickadee (*Poecile hudsonicus*) Conservation Status Rank Summary

December 12, 2024

For details on assessment and ranking methodology, see: [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#)

Rarity and Trends

Rank Factor	Date Assessed	Value	Score	Data Source	Comments
Rarity					
Range Extent	2024-12-12	Y: 52270.2 km ²	3.930	MTNHP Range Maps	None
Area of Occupancy	2024-12-12	2132 4km ² cells	4.130	MTNHP Modeling	None
Number of Occurrences	2024-12-12	92	4.130	MTNHP Databases	None
Population Size			-		Factor not used in ranking.
# of Occurrences in Good Condition			-		Factor not used in ranking.
% of Area Occupied in Good Condition			-		Factor not used in ranking.
Environmental Specificity	2009-01-29	Narrow	-	MTNHP Species Rank Data Table	Factor not used in ranking. Dependent on spruce-fir forest. Methodology: NS (2003) Original Score: B
Rarity is calculated by averaging weighted factor scores: $((3.93 \times 1) + (4.13 \times 2) + (4.13 \times 1)) / 4 = 4.08$					
Trends					
Short-term Trend	2024-12-12		0.000		None
Long-term Trend	2009-01-29		0.000	MTNHP Species Rank Data Table	Spruce-fir forest relatively stable since European arrival. Methodology: NS (2003) Original Score: E
Trends score is calculated by summing weighted short and long-term trend scores: $((0.00 \times 2) + (0.00 \times 1)) = 0.00$					

Threats

Rank Factor	Date Assessed	Value	Score	Data Source	Comments
Threats					
Overall Threat Impact		Very high - high	[0.000, 1.830]		Loss of mature or late successional forest may represent a threat to the species through increased fire, timber harvest, and insect outbreak since they are dependent on trees with soft heartwood (usually dead) for excavating nest holes.
Intrinsic Vulnerability	2009-01-29	Not intrinsically vulnerable	-	MTNHP Species Rank Data Table	Factor not used in ranking. Methodology: NS (2003) Original Score: C
Threat score is calculated from Overall Threat Impact when available or Intrinsic Vulnerability if not: ([0.00, 1.83]) = [0.00, 1.83]					

Individual Threats Data

Threat Category	Date Assessed	Impact Score	Scope	Severity	Immediacy	Comments
Biological Resource Use	2024-12-12	Medium	Restricted	Serious	High	Salvage logging of burned areas appears to have significant impacts on the species.
Natural System Modifications	2024-12-12	Medium - Low	Large	Moderate-Slight	High	Species is was detected in burned forest at approximate levels as unburned forest in one study, but author postulated this was due to the presence of trees that had survived the fire. Severe burns that resulted in loss of crowns would be expected to cause high impacts to the species.
Climate Change & Severe Weather	2024-12-12	High	Pervasive	Serious	Moderate	Audubon's Survival by Degrees project predicts a drastic reduction in species breeding habitat within Montana with an increase in temperatures of 1.5C.
Threat Tally: 0 - Very High, 1 - High, [1,2] - Medium, [0,1] - Low Overall Threat Impact* = Very high - high						

*See [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#) for calculation of Overall Threat Impact based on the number and impact of individual threats.

Conservation Status Rank Calculation

Raw score

Rarity: $(4.08 \times 70\%)$ + Threats: $([0.00, 1.83] \times 30\%)$ + Trends: $(0.00) = [2.86, 3.41]$

Calculated Rank: S3

Accepted Rank	S3B
Date Approved	2001-08-01
Approval Authority	Montana Species of Concern Committee
Rank Justification	Species is a relatively rare resident of coniferous forests in northwestern Montana. Populations appear to continue to occupy historic sites. It faces threats from a warming climate, salvage logging of burned areas, and high severity fires.

Supplementary Information

Montana Natural Heritage Program. 2021. Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species. 18 p.

https://mtnhp.mt.gov/docs/Montana_State_Rank_Criteria_20211201.pdf

Montana Field Guide Species Account:

<https://fieldguide.mt.gov/speciesDetail.aspx?elcode=ABPAW01060>

Predicted Suitable Habitat Model:

<https://mtnhp.mt.gov/resources/models/?elcode=ABPAW01060>

Information Needs

Information needs are assessed by considering the availability of factors used to assess species status as well as the quality of these assessments. Current information availability and quality to inform Conservation Status Rank for this species are highlighted.

Rank Factor	Assessment Category	Value	Criteria
General Status	Status Quality	Adequate	Calculated rank has low uncertainty and is represented by a single rank (e.g. S3); accepted rank may be adjusted to a range rank (e.g. S2S3)
		Poor	Rank assessed as SU or calculated rank has notable uncertainty and corresponds to a range rank with 2 or more values (e.g. S2?, S1S3, or S4S5)
Rarity	Range Quality	Adequate	Range polygon adequately represents area of probable occupancy and does not include substantial unoccupied areas; range may be adequately defined and still include areas of unsuitable habitat (e.g. mountain ranges for plains species)
		Marginal	Range polygon defined, but may include or exclude notable areas where the species may or may not occur on the landscape
		Poor	Range polygon not defined
	Habitat Quality	Adequate	Species-habitat relationship is well-defined (e.g. relevant literature or robust habitat model available)
		Marginal	Understanding of species-habitat relationship is adequate among some but not all habitats (e.g. literature covers similar habitats outside of Montana or habitat model performance is only somewhat adequate)
		Poor	Species-habitat relationship is not well understood
Threats	Threat Quality	Adequate	Threat Impact is a single value (including "Unthreatened")
		Marginal	Threat Impact assessed at more than one value (e.g. "High - Medium")
		Poor	Threat Impact is Unknown but Intrinsic Vulnerability is assessed
		Unknown	Threat Impact is Unknown and Intrinsic Vulnerability is not assessed
Trends	Recency	Current	Short-term Trend assessment date less than 10 years old
		Out of Date but Adequate	Short-term Trend assessment date is more than 10 years old or Unknown, but species is Unthreatened
		Out of Date	Short-term Trend assessment date more than 10 years old
		Not Available	Short-term Trend data are not available
	Trend Quality	Sufficient	Short-term Trend assessed at a single value or multiple values with a minimum trend greater than -10% (stable or increasing)
		Unknown but Sufficient	Short-term Trend is Unknown, but species is Unthreatened
		Poor	Short-term Trend is less than -10% (in decline) with two or more values selected
		Unknown	Short-term Trend is Unknown

Summary of Information Availability

Rarity data are available. Threat data are mostly inferred from studies in the boreal forests. Trend is based on continued incidental observations of the species within the historic breeding range.

Summary of Information Needs

General avian monitoring programs are insufficient to characterize population changes for this species. Species specific monitoring is needed to determine population trend and explore impacts of threats.

Additional Threat Details

The table below contains the complete threats assessment for this species. While the Conservation Status Rank Calculation is based on cumulative, broadly categorized (Level 1) threats data, threats are assessed and tracked for more specifically categorized (Level 2) threats when available.

Threat Category	Date Assessed	Assessed By	Data Source	Scope	Severity	Immediacy	Comments
Biological Resource Use - 5.3 - Logging & Wood Harvesting	2024-12-12	Dan Bachen	Hannon and Drapeau 2005	Restricted	Serious	High	Salvage logging of burned areas appears to have significant impacts on the species.
Natural System Modifications - 7.1 - Fire & Fire Suppression	2024-12-12	Dan Bachen	Morrisette 2000; Expert Opinion	Large	Moderate-Slight	High	Species is was detected in burned forest at approxiamte levels as unburned forest in one study, but author postulated this was due to the presence of trees that had survived the fire. Severe burns that resulted in loss of crowns would be expected to cause high impacts to the species.
Climate Change & Severe Weather - 11.1 - Habitat Shifting & Alteration	2024-12-12	Dan Bachen	Audubon Survival by Degrees project	Pervasive	Serious	Moderate	Audubon's Survival by Degrees project predicts a drastic reduction in species breeding habitat within Montana with an increase in temperatures of 1.5C.