

Marbled Godwit (*Limosa fedoa*)

Conservation Status Rank Summary

October 16, 2025

For details on assessment and ranking methodology, see: [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#)

Rarity and Trends

Rank Factor	Date Assessed	Value	Score	Data Source	Comments
Rarity					
Range Extent	2025-10-16	S: 201562.3 km ²	4.710	MTNHP Range Maps	None
Area of Occupancy	2025-10-16	11247 4km ² cells	4.810	MTNHP Modeling	None
Number of Occurrences			-		Factor not used in ranking.
Population Size			-		Factor not used in ranking.
# of Occurrences in Good Condition			-		Factor not used in ranking.
% of Area Occupied in Good Condition			-		Factor not used in ranking.
Environmental Specificity	2008-09-15	Moderate	-	MTNHP Species Rank Data Table	Factor not used in ranking. Dependent on prairie grasslands Methodology: NS (2003) Original Score: C
Rarity is calculated by averaging weighted factor scores: $((4.71 \times 1) + (4.81 \times 2)) / 3 = 4.78$					
Trends					
Short-term Trend	2023-12-20	[-0.6, 14.3%]	[0.000, 0.070]	IMBCR	IMBCR trend in population estimates for Montana. "- 95% CI"
Long-term Trend	2008-09-15		-0.140	MTNHP Species Rank Data Table	Very large reduction in prairie grassland cover types since the arrival of Europeans. Methodology: NS (2003) Original Score: D
Trends score is calculated by summing weighted short and long-term trend scores: $(((0.00, 0.07) \times 2) + (-0.14 \times 1)) = [-0.14, 0.00]$					

Threats

Rank Factor	Date Assessed	Value	Score	Data Source	Comments
Threats					
Overall Threat Impact		High	1.830		Conversion of prairie habitats
Intrinsic Vulnerability	2008-09-15	Not intrinsically vulnerable	-	MTNHP Species Rank Data Table	Factor not used in ranking. Methodology: NS (2003) Original Score: C
Threat score is calculated from Overall Threat Impact when available or Intrinsic Vulnerability if not: (1.83) = 1.83					

Individual Threats Data

Threat Category	Date Assessed	Impact Score	Scope	Severity	Immediacy	Comments
Agriculture & Aquaculture	2025-10-16	Medium	Large	Moderate	High	Loss of native grassland habitat to conversion to row crop agriculture.
Natural System Modifications	2026-03-19	Medium	Large	Moderate	High	In northern prairies of Canada and U.S., breeds in short, sparsely to moderately vegetated landscapes that include native grassland and wetland complexes with a variety of wetland classes (ephemeral to semipermanent; Johnson et al. 1998). Draining and plowing of temporary and ephemeral wetlands for crops and maintenance of high water levels for tourism and irrigation also are problems. Managed wetlands can be advantageous if they have shallow water and little vegetation at edges (see Habitat, above), not steep slopes or cattail. In wetland complexes advantageous if variety of water levels present, including very shallow wetlands.
Climate Change & Severe Weather	2026-03-19	Medium	Pervasive	Moderate	Moderate	Audubon's climate model predicts that as temperatures increase, Marbled Godwit will suffer from impacts of drought.
Threat Tally: 0 - Very High, 0 - High, 3 - Medium, 0 - Low Overall Threat Impact* = High						

*See [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#) for calculation of Overall Threat Impact based on the number and impact of individual threats.

Conservation Status Rank Calculation

Raw score

Rarity: $(4.78 \times 70\%)$ + Threats: $(1.83 \times 30\%)$ + Trends: $([-0.14, 0.00])$ = $[3.75, 3.89]$

Calculated Rank: S4

Accepted Rank	S4B
Date Approved	Date Unknown
Approval Authority	Legacy Assessment: MTNHP Staff
Rank Justification	Species is found in steppe habitat across central and eastern Montana. Populations appear stable. Threats include loss of native habitat through conversion to row crop agriculture.

Supplementary Information

Montana Natural Heritage Program. 2021. Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species. 18 p.

https://mtnhp.mt.gov/docs/Montana_State_Rank_Criteria_20211201.pdf

Montana Field Guide Species Account:

<https://fieldguide.mt.gov/speciesDetail.aspx?elcode=ABNNF08040>

Predicted Suitable Habitat Model:

<https://mtnhp.mt.gov/resources/models/?elcode=ABNNF08040>

Information Needs

Information needs are assessed by considering the availability of factors used to assess species status as well as the quality of these assessments. Current information availability and quality to inform Conservation Status Rank for this species are highlighted.

Rank Factor	Assessment Category	Value	Criteria
General Status	Status Quality	Adequate	Calculated rank has low uncertainty and is represented by a single rank (e.g. S3); accepted rank may be adjusted to a range rank (e.g. S2S3)
		Poor	Rank assessed as SU or calculated rank has notable uncertainty and corresponds to a range rank with 2 or more values (e.g. S2?, S1S3, or S4S5)
Rarity	Range Quality	Adequate	Range polygon adequately represents area of probable occupancy and does not include substantial unoccupied areas; range may be adequately defined and still include areas of unsuitable habitat (e.g. mountain ranges for plains species)
		Marginal	Range polygon defined, but may include or exclude notable areas where the species may or may not occur on the landscape
		Poor	Range polygon not defined
	Habitat Quality	Adequate	Species-habitat relationship is well-defined (e.g. relevant literature or robust habitat model available)
		Marginal	Understanding of species-habitat relationship is adequate among some but not all habitats (e.g. literature covers similar habitats outside of Montana or habitat model performance is only somewhat adequate)
		Poor	Species-habitat relationship is not well understood
Threats	Threat Quality	Adequate	Threat Impact is a single value (including "Unthreatened")
		Marginal	Threat Impact assessed at more than one value (e.g. "High - Medium")
		Poor	Threat Impact is Unknown but Intrinsic Vulnerability is assessed
		Unknown	Threat Impact is Unknown and Intrinsic Vulnerability is not assessed
Trends	Recency	Current	Short-term Trend assessment date less than 10 years old
		Out of Date but Adequate	Short-term Trend assessment date is more than 10 years old or Unknown, but species is Unthreatened
		Out of Date	Short-term Trend assessment date more than 10 years old
		Not Available	Short-term Trend data are not available
	Trend Quality	Sufficient	Short-term Trend assessed at a single value or multiple values with a minimum trend greater than -10% (stable or increasing)
		Unknown but Sufficient	Short-term Trend is Unknown, but species is Unthreatened
		Poor	Short-term Trend is less than -10% (in decline) with two or more values selected
		Unknown	Short-term Trend is Unknown

Summary of Information Availability

Information to assess status is available.

Summary of Information Needs

No additional information is needed.

Additional Threat Details

The table below contains the complete threats assessment for this species. While the Conservation Status Rank Calculation is based on cumulative, broadly categorized (Level 1) threats data, threats are assessed and tracked for more specifically categorized (Level 2) threats when available.

Threat Category	Date Assessed	Assessed By	Data Source	Scope	Severity	Immediacy	Comments
Agriculture & Aquaculture - 2.1 - Annual & Perennial Non-Timber Crops	2025-10-16	Dan Bachen	Expert Opinion	Large	Moderate	High	Loss of native grassland habitat to conversion to row crop agriculture.
Natural System Modifications - 7.2 - Dams & Water Management/Use	2026-03-19	Dan Bachen	SWAP Assessment	Large	Moderate	High	In northern prairies of Canada and U.S., breeds in short, sparsely to moderately vegetated landscapes that include native grassland and wetland complexes with a variety of wetland classes (ephemeral to semipermanent; Johnson et al. 1998). Draining and plowing of temporary and ephemeral wetlands for crops and maintenance of high water levels for tourism and irrigation also are problems. Managed wetlands can be advantageous if they have shallow water and little vegetation at edges (see Habitat, above), not steep slopes or cattail. In wetland complexes advantageous if variety of water levels present, including very shallow wetlands.
Invasive & Other Problematic Species, Genes & Diseases - 8.4 - Problematic Species/Diseases of Unknown Origin	2026-03-19	Dan Bachen	SWAP Assessment	Unknown	Unknown	High	Staging fall migrants have been found dead in avian botulism outbreaks (bacterium Clostridium botulinum type C) in s. Canadian Prairies.
Climate Change & Severe Weather - 11.2 - Droughts	2026-03-19	Dan Bachen	SWAP Assessment	Pervasive	Moderate	Moderate	Audubon's climate model predicts that as temperatures increase, Marbled Godwit will suffer from impacts of drought.