

Turkey Vulture (*Cathartes aura*)

Conservation Status Rank Summary

February 6, 2025

For details on assessment and ranking methodology, see: [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#)

Rarity and Trends

| Rank Factor | Date Assessed | Value | Score | Data Source | Comments |
|--|---------------|--------------------------------|-------|-------------------------------|--|
| Rarity | | | | | |
| Range Extent | 2025-02-06 | S: 380530.8 km ² | 4.710 | MTNHP Range Maps | None |
| Area of Occupancy | 2025-02-06 | 10736 4km ² cells | 4.810 | MTNHP Modeling | None |
| Number of Occurrences | | | - | | Factor not used in ranking. |
| Population Size | | | - | | Factor not used in ranking. |
| # of Occurrences in Good Condition | | | - | | Factor not used in ranking. |
| % of Area Occupied in Good Condition | | | - | | Factor not used in ranking. |
| Environmental Specificity | 2011-12-22 | Moderate | - | MTNHP Species Rank Data Table | Factor not used in ranking. Moderate Generalist. Forage over a large variety of habitats, but usually nest in caves which are relatively rare on the landscape. Methodology: NS (2003) Original Score: C |
| Rarity is calculated by averaging weighted factor scores: $((4.71 \times 1) + (4.81 \times 2)) / 3 = 4.78$ | | | | | |
| Trends | | | | | |
| Short-term Trend | 2023-12-20 | 20.2% | 0.070 | IMBCR | IMBCR trend in population estimates for Montana. "-Point Estimate" |
| Long-term Trend | 2011-12-22 | | 0.000 | MTNHP Species Rank Data Table | Populations likely declined after loss of Bison, but have likely increased to within +/- 25% of those pre-European levels. Methodology: NS (2003) Original Score: E |
| Trends score is calculated by summing weighted short and long-term trend scores: $((0.07 \times 2) + (0.00 \times 1)) = 0.14$ | | | | | |

Threats

| Rank Factor | Date Assessed | Value | Score | Data Source | Comments |
|--|---------------|-----------------------|-------|-------------------------------|--|
| Threats | | | | | |
| Overall Threat Impact | | Low/No Threats | 5.500 | | Contaminants such as lead from gut piles and human persecution probably represent the greatest threats to the species. |
| Intrinsic Vulnerability | 2011-12-22 | Moderately vulnerable | - | MTNHP Species Rank Data Table | Factor not used in ranking. Moderately Vulnerable. Species exhibits moderate age of maturity, frequency of reproduction, and/or fecundity such that populations generally tend to recover from decreases in abundance within 5-20 years or 2-5 generations. Species has good dispersal capabilities such that extirpated populations generally become reestablished through natural recolonization. Methodology: NS (2003) Original Score: B |
| Threat score is calculated from Overall Threat Impact when available or Intrinsic Vulnerability if not: (5.50) = 5.50 | | | | | |

Individual Threats Data

| Threat Category | Date Assessed | Impact Score | Scope | Severity | Immediacy | Comments |
|---|---------------|--------------|-----------|----------|-----------|---|
| Biological Resource Use | 2025-02-06 | Low | Pervasive | Slight | High | Lead poisoning from fragmented bullets in carcasses may represent a threat to the species as it does other scavengers. Due to the species migratory nature of the species almost all individuals have left Montana by the time general rifle season opens in late October reducing or eliminating this threat in the state. |
| Threat Tally: 0 - Very High, 0 - High, 0 - Medium, 1 - Low Overall Threat Impact* = Low/No Threats | | | | | | |

*See [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#) for calculation of Overall Threat Impact based on the number and impact of individual threats.

Conservation Status Rank Calculation

Raw score

Rarity: $(4.78 \times 70\%)$ + Threats: $(5.50 \times 30\%)$ + Trends: (0.14) = 5.13

Calculated Rank: S5

| | |
|---------------------------|---|
| Accepted Rank | S5B |
| Date Approved | 2025-02-06 |
| Approval Authority | Montana Natural Heritage Program Staff |
| Rank Justification | Species is relatively common within suitable habitat and widely distributed across portions of the state. |

Supplementary Information

Montana Natural Heritage Program. 2021. Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species. 18 p.

https://mtnhp.mt.gov/docs/Montana_State_Rank_Criteria_20211201.pdf

Montana Field Guide Species Account:

<https://fieldguide.mt.gov/speciesDetail.aspx?elcode=ABNKA02010>

Predicted Suitable Habitat Model:

<https://mtnhp.mt.gov/resources/models/?elcode=ABNKA02010>

Information Needs

Information needs are assessed by considering the availability of factors used to assess species status as well as the quality of these assessments. Current information availability and quality to inform Conservation Status Rank for this species are highlighted.

| Rank Factor | Assessment Category | Value | Criteria |
|----------------|---------------------|--------------------------|---|
| General Status | Status Quality | Adequate | Calculated rank has low uncertainty and is represented by a single rank (e.g. S3); accepted rank may be adjusted to a range rank (e.g. S2S3) |
| | | Poor | Rank assessed as SU or calculated rank has notable uncertainty and corresponds to a range rank with 2 or more values (e.g. S2?, S1S3, or S4S5) |
| Rarity | Range Quality | Adequate | Range polygon adequately represents area of probable occupancy and does not include substantial unoccupied areas; range may be adequately defined and still include areas of unsuitable habitat (e.g. mountain ranges for plains species) |
| | | Marginal | Range polygon defined, but may include or exclude notable areas where the species may or may not occur on the landscape |
| | | Poor | Range polygon not defined |
| | Habitat Quality | Adequate | Species-habitat relationship is well-defined (e.g. relevant literature or robust habitat model available) |
| | | Marginal | Understanding of species-habitat relationship is adequate among some but not all habitats (e.g. literature covers similar habitats outside of Montana or habitat model performance is only somewhat adequate) |
| | | Poor | Species-habitat relationship is not well understood |
| Threats | Threat Quality | Adequate | Threat Impact is a single value (including "Unthreatened") |
| | | Marginal | Threat Impact assessed at more than one value (e.g. "High - Medium") |
| | | Poor | Threat Impact is Unknown but Intrinsic Vulnerability is assessed |
| | | Unknown | Threat Impact is Unknown and Intrinsic Vulnerability is not assessed |
| Trends | Recency | Current | Short-term Trend assessment date less than 10 years old |
| | | Out of Date but Adequate | Short-term Trend assessment date is more than 10 years old or Unknown, but species is Unthreatened |
| | | Out of Date | Short-term Trend assessment date more than 10 years old |
| | | Not Available | Short-term Trend data are not available |
| | Trend Quality | Sufficient | Short-term Trend assessed at a single value or multiple values with a minimum trend greater than -10% (stable or increasing) |
| | | Unknown but Sufficient | Short-term Trend is Unknown, but species is Unthreatened |
| | | Poor | Short-term Trend is less than -10% (in decline) with two or more values selected |
| | | Unknown | Short-term Trend is Unknown |

Summary of Information Availability

Information to assess status is available.

Summary of Information Needs

No further information is needed.

Additional Threat Details

The table below contains the complete threats assessment for this species. While the Conservation Status Rank Calculation is based on cumulative, broadly categorized (Level 1) threats data, threats are assessed and tracked for more specifically categorized (Level 2) threats when available.

| Threat Category | Date Assessed | Assessed By | Data Source | Scope | Severity | Immediacy | Comments |
|--|---------------|-------------|-------------------|-----------|----------|-----------|---|
| Biological Resource Use - 5.1 - Hunting & Collecting Terrestrial Animals | 2025-02-06 | Dan Bachen | Expert Opinion | Pervasive | Slight | High | Lead poisoning from fragmented bullets in carcasses may represent a threat to the species as it does other scavengers. Due to the species migratory nature of the species almost all individuals have left Montana by the time general rifle season opens in late October reducing or eliminating this threat in the state. |