

White-winged Crossbill (*Loxia leucoptera*)

Conservation Status Rank Summary

January 30, 2025

For details on assessment and ranking methodology, see: [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#)

Rarity and Trends

Rank Factor	Date Assessed	Value	Score	Data Source	Comments
Rarity					
Range Extent	2025-01-30	Y: 144219.0 km ²	3.930	MTNHP Range Maps	None
Area of Occupancy	2025-01-30	17543 4km ² cells	5.500	MTNHP Modeling	None
Number of Occurrences			-		Factor not used in ranking.
Population Size			-		Factor not used in ranking.
# of Occurrences in Good Condition			-		Factor not used in ranking.
% of Area Occupied in Good Condition			-		Factor not used in ranking.
Environmental Specificity	2011-12-21	Narrow	-	MTNHP Species Rank Data Table	Factor not used in ranking. Narrow specialist. Species has morphologically specialized feeding adaptations and favors late successional mesic conifer forests with larch and spruce seeds which are their preferred diet items. Methodology: NS (2003) Original Score: B
Rarity is calculated by averaging weighted factor scores: $((3.93 \times 1) + (5.50 \times 2)) / 3 = 4.98$					
Trends					
Short-term Trend	2023-12-20	-14.9%	-0.070	IMBCR	IMBCR trend in population estimates for Montana. "-Point Estimate"
Long-term Trend	2011-12-21		0.000	MTNHP Species Rank Data Table	Species favors mesic conifer forests with larch and spruce seeds as their favored diet items. Mesic conifer forests in the Northern Rockies have remained relatively stable within +/- 25% of pre European levels. Methodology: NS (2003) Original Score: E
Trends score is calculated by summing weighted short and long-term trend scores: $((-0.07 \times 2) + (0.00 \times 1)) = -0.14$					

Threats

Rank Factor	Date Assessed	Value	Score	Data Source	Comments
Threats					
Overall Threat Impact		Medium	3.670		Loss of mature or late successional forest may represent a threat to the species through increased fire, timber harvest, and insect outbreak since they tend to favor foraging in mature forests. Species is irruptive in association with cone crops.
Intrinsic Vulnerability	2011-12-21	Not intrinsically vulnerable	-	MTNHP Species Rank Data Table	Factor not used in ranking. Not Intrinsically Vulnerable. Species matures quickly, reproduces frequently, and/or has a high fecundity such that populations recover quickly (5 years or 2 generations) from decreases in abundance. Species has good dispersal capabilities such that extirpated populations generally become reestablished through natural recolonization. Methodology: NS (2003) Original Score: C
Threat score is calculated from Overall Threat Impact when available or Intrinsic Vulnerability if not: (3.67) = 3.67					

Individual Threats Data

Threat Category	Date Assessed	Impact Score	Scope	Severity	Immediacy	Comments
Biological Resource Use	2025-01-30	Low	Restricted	Moderate	High	Loss of mature forest and fragmentation due to logging
Natural System Modifications	2025-01-30	Medium	Large	Moderate	High	Loss of mature forest and fragmentation due to fire and fire suppression
Threat Tally: 0 - Very High, 0 - High, 1 - Medium, 1 - Low Overall Threat Impact* = Medium						

*See [Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species](#) for calculation of Overall Threat Impact based on the number and impact of individual threats.

Conservation Status Rank Calculation

Raw score

Rarity: $(4.98 \times 70\%)$ + Threats: $(3.67 \times 30\%)$ + Trends: (-0.14) = 4.45

Calculated Rank: S4

Accepted Rank	S4B
Date Approved	2025-01-30
Approval Authority	MTNHP Staff
Rank Justification	

Supplementary Information

Montana Natural Heritage Program. 2021. Conservation Status Assessment Definitions, Process, Rank Factors, and Calculation of State Ranks for Montana Species. 18 p.

https://mtnhp.mt.gov/docs/Montana_State_Rank_Criteria_20211201.pdf

Montana Field Guide Species Account:

<https://fieldguide.mt.gov/speciesDetail.aspx?elcode=ABPBY05020>

Predicted Suitable Habitat Model:

<https://mtnhp.mt.gov/resources/models/?elcode=ABPBY05020>

Information Needs

Information needs are assessed by considering the availability of factors used to assess species status as well as the quality of these assessments. Current information availability and quality to inform Conservation Status Rank for this species are highlighted.

Rank Factor	Assessment Category	Value	Criteria
General Status	Status Quality	Adequate	Calculated rank has low uncertainty and is represented by a single rank (e.g. S3); accepted rank may be adjusted to a range rank (e.g. S2S3)
		Poor	Rank assessed as SU or calculated rank has notable uncertainty and corresponds to a range rank with 2 or more values (e.g. S2?, S1S3, or S4S5)
Rarity	Range Quality	Adequate	Range polygon adequately represents area of probable occupancy and does not include substantial unoccupied areas; range may be adequately defined and still include areas of unsuitable habitat (e.g. mountain ranges for plains species)
		Marginal	Range polygon defined, but may include or exclude notable areas where the species may or may not occur on the landscape
		Poor	Range polygon not defined
	Habitat Quality	Adequate	Species-habitat relationship is well-defined (e.g. relevant literature or robust habitat model available)
		Marginal	Understanding of species-habitat relationship is adequate among some but not all habitats (e.g. literature covers similar habitats outside of Montana or habitat model performance is only somewhat adequate)
		Poor	Species-habitat relationship is not well understood
Threats	Threat Quality	Adequate	Threat Impact is a single value (including "Unthreatened")
		Marginal	Threat Impact assessed at more than one value (e.g. "High - Medium")
		Poor	Threat Impact is Unknown but Intrinsic Vulnerability is assessed
		Unknown	Threat Impact is Unknown and Intrinsic Vulnerability is not assessed
Trends	Recency	Current	Short-term Trend assessment date less than 10 years old
		Out of Date but Adequate	Short-term Trend assessment date is more than 10 years old or Unknown, but species is Unthreatened
		Out of Date	Short-term Trend assessment date more than 10 years old
		Not Available	Short-term Trend data are not available
	Trend Quality	Sufficient	Short-term Trend assessed at a single value or multiple values with a minimum trend greater than -10% (stable or increasing)
		Unknown but Sufficient	Short-term Trend is Unknown, but species is Unthreatened
		Poor	Short-term Trend is less than -10% (in decline) with two or more values selected
		Unknown	Short-term Trend is Unknown

Summary of Information Availability

None

Summary of Information Needs

None

Additional Threat Details

The table below contains the complete threats assessment for this species. While the Conservation Status Rank Calculation is based on cumulative, broadly categorized (Level 1) threats data, threats are assessed and tracked for more specifically categorized (Level 2) threats when available.

Threat Category	Date Assessed	Assessed By	Data Source	Scope	Severity	Immediacy	Comments
Biological Resource Use - 5.3 - Logging & Wood Harvesting	2025-01-30	Dan Bachen	Expert Opinion	Restricted	Moderate	High	Loss of mature forest and fragmentation due to logging
Natural System Modifications - 7.1 - Fire & Fire Suppression	2025-01-30	Dan Bachen	Expert Opinion	Large	Moderate	High	Loss of mature forest and fragmentation due to fire and fire suppression